Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

UNITED STATES DEPARTMENT OF AGRICULTURE
Miscellaneous Publication No. 372

Washington, D. C. May 1940

LIBRARY
JUL 9 - 1940
AGRICULTURAL ECONOMICS
U. S. DEPARTMENT OF AGRICULTURE

A SURVEY AND DISCUSSION OF LYSIMETERS

AND

A BIBLIOGRAPHY ON THEIR CONSTRUCTION AND PERFORMANCE

By

HELMUT KOHNKE and F. R. DREIBELBIS
Assistant Soil Technologists, Hydrologic Division

and

J. M. DAVIDSON
Assistant Soil Conservationist
Conservation Experiment Stations Division
Soil Conservation Service

For sale by the Superintendent of Documents, Washington, D. C. - - - - - - - - - - Price 10 cents

UNITED STATES
DEPARTMENT OF AGRICULTURE
MISCELLANEOUS PUBLICATION NO. 372

Washington, D. C. May 1940

A SURVEY AND DISCUSSION OF LYSIMETERS AND A BIBLIOGRAPHY ON THEIR CONSTRUCTION AND PERFORMANCE

By Helmut Kohnke and F. R. Dreibelbis, *assistant soil technologists, Hydrologic Division,* and J. M. Davidson, *assistant soil conservationist, Conservation Experiment Stations Division, Soil Conservation Service*

CONTENTS

	Page
Lysimeters: A survey and discussion	1
Introduction	1
Classification of lysimeters	2
History of lysimeter studies	5
Discussion	19
Established lysimeter findings	24
Findings with respect to the moisture relationships of soils	24
Findings with respect to the chemical constituents of the percolate	25
Suggestions for future lysimeter studies	27
Construction of lysimeters	27
Experimental plan	29
Publication of data	30
Supplementary studies	30
Lysimeters: A survey and discussion—Con.	
Conclusion	30
Bibliography on lysimeters	31
Content and organization	31
Bibliography by countries	32
United States	32
Great Britain	45
Asia	50
Europe	52
Africa	63
Author index	65

LYSIMETERS: A SURVEY AND DISCUSSION

By Helmut Kohnke and F. R. Dreibelbis [1]

INTRODUCTION

With the great impetus that soil conservation and flood-control work have given to the study of the relations of water to soil, lysimeters have received wider attention during the past decade than at any previous time. They have been used for at least two and a half centuries in studies of the percolation of water through soil. The purpose of such studies may be strictly hydrological, attention being given only to the rate and amount of the percolate. In most of the more recent investigations the chemical composition of the percolate is analyzed, generally as a part of the study of the fertility balance of the soil, but, in some cases, in connection with studies of the genesis and development of soils.

[1] The authors wish to express their appreciation to W. U. Garstka, Hydrologic Division, Soil Conservation Service, for guidance and valuable assistance given in the development of this publication.

CLASSIFICATION OF LYSIMETERS

Lysimeters, classified according to the principles of construction, are of three major types: (1) Monolith, or undisturbed soil-block; (2) Ebermayer, and (3) Filled-in.

In the first type, a case is built around the sides of a block of soil as it is found in the field, a partly open bottom is attached, and the percolate is conducted to receiver tanks. In the Ebermayer type, the soil is left in situ and a percolate collecting funnel is placed under it, but no side walls separate a definite soil block from the adjoining soil. A tube attached to the funnel conveys the percolate into a receptacle. The filled-in type consists of a container, which has vertical walls, an open top, and a bottom that provides for percolation and is filled with soil that has been removed from its original location and that usually has been screened and mixed in order to make it uniform. In some lysimeters of this type the tops of the side walls are completely covered with soil so that the ground is level with the surrounding soil. This construction permits natural run-off and eliminates the border effect resulting from the unplanted area along the rim of the lysimeter. In these lysimeters an attempt is made to combine the advantages of both the filled-in and the Ebermayer types. A special kind of the filled-in type can be called the laboratory lysimeter, which is usually small in size and artificially supplied with water or a solution. It is used to study specific problems; and experiments with it are usually restricted to short periods because the rather unnatural conditions tend quickly to deteriorate the soil.

Lysimeters may also be classified according to the type of provision made for the run-off of rain water that falls in excess of the infiltration rate. The following classes are in use:

(1) Unlimited run-off: The Ebermayer lysimeters allow unrestricted run-off, since they have no side walls. A number of the more recent lysimeters of the soil-block type (table 1) are constructed to permit run-off, as are also the filled-in types at San Dimas and Berkeley, Calif. (*4, 8, 10*),[2] and at La Crosse, Wis. (*205*).

(2) Run-off through overflow pipes: Some lysimeters are provided with overflow pipes a short distance above the soil surface. This permits the water of very large rainstorms to run off, but a certain amount of water will remain standing on the soil surface. Dalton (*211*) recognized the necessity of natural run-off and installed a vertical overflow pipe in his first lysimeter. The soil soon settled below the level of the pipe opening, however, making the device ineffective. Among the more recent installations only the two sets of lysimeters in India are equipped with horizontal overflow pipes, which are about 3 inches above the ground.

(3) No run-off: The great majority of the lysimeters of the filled-in type (and a number of the soil-block type) are encased with walls that extend above the soil surface on all sides. Such lysimeters dispose of the rainfall entirely through percolation, evaporation, or transpiration. The absence of provision for outflow of run-off results in a stagnation of water on the top of the soil after nearly every rain, particularly on heavy soils.

[2] Italic numbers in parentheses refer to Bibliography, p. 32.

TABLE 1.—*Data on soil-block and Ebermayer lysimeters with undisturbed soils*

SOIL-BLOCK LYSIMETERS

Investigator and citation number in bibliography, or footnote	Location	Year of installation	Precipitation	Drainage as percentage of precipitation	Provision for run-off	Remarks
			Inches	*Percent*		
Lawes and Gilbert (early work) Miller (later work) (*224, 227, 242, 252*).	Rothamsted, England.	1870	29.0	47.1 to 53.2	No	
E. L. Sturtevant (*77, 84, 111*).	Massachusetts	1875			No	
Sturtevant and Babcock (*93, 122*).	Geneva, N. Y	1882		[1] 37.6 [2] 41.5 [3] 43.8	No	
Babcock and Goff (*99*)	do	1887			No	
J. W. Sanborn (*85*)	Columbia, Mo	1888			No	
Burt and Leather (*290*)	Cawnpore, India	1903	31.5	44.3 to 48.9	Yes	
R. E. Horton (*103*)	Grafenburg, N. Y	1904			No	
Welbel (*463, 474, 476*)	Ploti, Soviet Union	1905	16.0	9.6 to 11.4	No	
Arnott and Leather (*284*)	Pusa, India	1906	40.0	25.0	Yes	
Hendrick and Welsh (*263, 267, 271, 272, 276, 278*).	Aberdeen, Scotland	1914	34.6	52 to 55	No	
Miller and Duley (*86*)	Columbia, Mo	1918			Yes	
Weaver and Crist [4]	Lincoln, Nebr	1923	[5]		Yes	With weighing mechanism.
Duley (*87*)	Columbia, Mo	1925			No	
Winnik (*308*)	Palestine	1925	20.0	17.0	No	
Harper [6]	Oklahoma	1927			Yes	
F. Shreve (*2*)	Tucson, Ariz	1929				
Musgrave (*65*)	Temple, Tex	1931			Yes	
California Forest and Range Experiment Station.[7]	Berkeley, Calif	1932			Yes	
Southwestern Forest and Range Experiment Station.[7]	Sierra Ancha Experimental Forest, Globe, Ariz.	1932			Yes	
Musgrave (*65, 69*)	Clarinda, Iowa	1933	[8] 25.6	[8] 5.6 to 22.9	Yes	With tensiometers for measuring capillary tension of soil moisture.
Lake States Forest Experiment Station (*205*).	Upper Mississippi Valley Soil and Water Conservation Experiment Station, La Crosse, Wis.	1934			Yes	
Southwestern Forest and Range Experiment Station.[7]	Sierra Ancha Experimental Forest, Globe, Ariz.	1934			Yes	
Musgrave [9]	Statesville, N. C	1934			Yes	
Southwestern Forest and Range Experiment Station.[7]	Fort Valley Branch Station, Fort Valley, Ariz.	1935			Yes	
Osugi (*313*)	Kyoto, Japan	1935 [10]			No	
Stauffer and Smith (*59*)	Urbana, Ill	1935			Yes	
North Appalachian Experimental Watershed, Soil Conservation Service (*138, 139, 140*).	Coshocton, Ohio	1936	[11] 45.0	[11] 60.5	Yes	With self-recording weighing mechanism.
Southwestern Forest and Range Experiment Station.[7]	Sierra Ancha Experimental Forest, Globe, Ariz.	1937			Yes	
Southwestern Forest and Range Experiment Station.[7]	do	1938			Yes	

[1] Surface of the lysimeter sodded.
[2] Surface of the lysimeter bare.
[3] Surface of the lysimeter cultivated.
[4] WEAVER, J. E., and CRIST, J. W. DIRECT MEASUREMENT OF WATER LOSS FROM VEGETATION WITHOUT DISTURBING THE NORMAL STRUCTURE OF THE SOIL. Ecology 5: 153–170. 1924.
[5] Irrigated.
[6] HARPER, H. J. A STUDY OF LYSIMETER INSTALLATIONS FOR THE MEASUREMENT OF GRAVITATIONAL WATER. Unpublished paper presented before the meeting of the Amer. Soc. Agron., November 1937.
[7] Unpublished data through courtesy of U. S. Forest Service.
[8] Musgrave and Neal (*68*).
[9] Private communication.
[10] Year published.
[11] Data for 1 year only.

TABLE 1.—*Data on soil-block and Ebermayer lysimeters with undisturbed soils*—Continued

EBERMAYER LYSIMETERS

Investigator and citation number in bibliography, or footnote	Location	Year of installation	Precipitation	Drainage as percentage of precipitation	Provision for run-off	Remarks
			Inches	*Percent*		
Welbel (*463, 474, 476*)	Ploti, Soviet Union	1903	16.0	1.9 to 11.4	Yes	
W. W. Gemmerling (*481*)	Moscow, Soviet Union.	1922			Yes	
V. P. Popov (*482*)	Soviet Union	1928			Yes	
J. S. Joffe (*88, 89, 90, 91*)	New Brunswick, N. J.	1929	35.8	.3 to 37.6	Yes	
R. C. Collison (*131*)	Geneva, N. Y.	1932		4.0 to 18.8	Yes	
H. A. Lunt (*19, 23*)	New Haven, Conn.	1933		11.1 to 85.8	Yes	
H. J. Harper [6]	Oklahoma				Yes	
University of Pretoria (*489*).	Pretoria, South Africa.	1934			Yes	

[6] HARPER, H. J. A STUDY OF LYSIMETRR INSTALLATIONS FOR THE MEASUREMENT OF GRAVITATIONAL WATER. Unpublished paper presented before the meeting of the Amer. Soc. Agron., November 1937.

Lysimeters may also be classified in two groups: Those that are and those that are not weighed. The very nature of the Ebermayer lysimeters prohibits any weighing, whereas both the soil-block and the filled-in types can be weighed. Because the weighing of a large mass of soil in a heavy container is a cumbersome task, not many lysimeters have been installed with weighing mechanisms. Investigators interested in studying only the fertility balance of soils can easily dispense with such installations. Some lysimeters (*351, 357*) [3] are so arranged that scales can be rolled underneath them. In only two cases, so far as the authors know, is a set of scales placed permanently under the lysimeter (*139, 140, 396, 397*).

In a great majority of experiments the moisture content of lysimeter soils has not been recorded. This is due to the difficulties connected with such determinations and also to the lack of appreciation of the importance of information on moisture content. A number of investigators determine the moisture content of the soil by weighing the whole lysimeter. This gives accurate data only where the dry weight of the original soil is available, as it may be in refilled lysimeters; for soil-block lysimeters the data will at best be approximate. The weighing method can give only the average moisture content of the total soil and does not indicate the distribution of the moisture. Weighing of lysimeters containing soil on top and sand on the bottom does not yield any direct data with respect to soil moisture. A similar objection can be raised against weighing soil-block lysimeters, since the natural soil profile tends to distribute soil moisture unevenly. Lysimeters filled with only one type of uniformly mixed soil may have different moisture contents at various depths. It should be kept in mind that the purpose of weighing lysimeters usually is not the determination of soil moisture but the tracing of evaporation and transpiration, for which purpose information regarding the distribution of soil moisture may not be necessary.

The direct determination of soil moisture in lysimeters might be carried out by removing soil samples and making a gravimetric determination. This technique, however, would destroy the lysimeter, since the openings left in the soil would permit excessive percolation. Under special circumstances destruction of the lysimeter would not be objectionable, as, for instance, when the exact soil moisture in a lysimeter at a definite time is the object of research. For the ordinary

[3] Personal communication from H. Baumann, concerning Berlin-Dahlem lysimeters, March 1938.

type of lysimeter installation a direct measurement of soil moisture throughout the profile without disturbing the soil should be considered a prerequisite. Tensiometers have recently been used successfully (*69*) for this purpose. The effective range of tensiometers in indicating soil moisture, however, is not wide enough for all lysimeter investigations. The development of a reliable in situ soil-moisture meter is urgently needed.

Two main types of lysimeters can be distinguished with respect to the method by which the percolating water is conducted from the soil to the receiver tanks. In one, the soil (or in the deeper soil-block type lysimeters, the parent material) rests immediately on the funnel or perforated bottom plate; in the other, a layer of sand or gravel is placed between the soil and the lysimeter bottom. The percolating water on its way to the receivers is always protected from evaporation, and in some lysimeters (*152*) it is protected also from oxidation.

HISTORY OF LYSIMETER STUDIES

It probably will never be known who was the first to study the percolation of water through soil, but it seems likely that such experiments were carried out by the natural philosophers of the Classical Age. The first actual lysimeter investigation, the report of which the writers have been able to study, was started in 1688 in Paris, by De la Hire [4] (*402*), who was interested in determining the origin of springs. At that time and for almost two centuries later the belief was held by some scientists that springs were entirely unrelated to rainfall, but originated from huge underground reservoirs that were replenished by the condensation of moist air or by ocean water entering through large crevasses. Others thought that springs represented that part of the precipitation that percolated through the soil and reappeared where a ground-water horizon cropped out. De la Hire used leaden vessels of three different depths (8 inches, 16 inches, and 8 feet), which he filled with sandy loam from the park of the castle of Louis XIV, whom he served as mathematician and meteorologist. He found that the lysimeters in grass evaporated more water than the ones in fallow, and he noticed a number of other phenomena that have been verified by many later lysimeter investigations. His findings with respect to the origin of springs, however, were not conclusive enough to convince the scientific world of that period.

Also of the filled-in type were the lysimeters of Maurice [5] (*223*), in Switzerland, and of Dalton (*211*), in England, both of whom started their experiments in 1796. Dalton, though commonly credited with being the first one to install lysimeters, acknowledges the inspiration he received from De la Hire.

In table 2 is given a record of lysimeter installations beginning with that of De la Hire in 1688 and continuing to those of the present. Data pertaining to the soil used, as well as to the type of construction, are given insofar as they are available in the literature. While an attempt was made to make a complete survey of the literature, it is most likely that there are some omissions. The historic references and discussions of findings are based entirely on the literature cited. The Forest Service supplied information concerning a number of their lysimeters for which no published references are available.

Table 3 gives the chronology of the development of lysimeters.

[4] De la Hire, Philippe. sur l'origine des rivières. Hist. de l'Acad. Roy. des Sci., pp. 1-6. 1703.
[5] Maurice. bibliographie universelle de genève sciences et arts. v. 1. No date. [Original not seen.]

TABLE 2.—*Lysimeter*

Investigator and citation number in bibliography, or footnote	Location	Year of installation	Soil	
			Texture	Other characteristics
De la Hire [1] (*402*)	Rungis, near Paris, France.	1688	Sand, loam	
John Dalton (*211*)	Manchester, England	1795	"Good fresh soil"	
Maurice [2]	Geneva, Switzerland	1796		
Gasparin [3]	Orange, France	1821		
Dickinson and Evans (*223*).	Nash Mills, England	1836	"Surface soil"	
John Dickinson (*214, 215*)	Abbottshill, England	1836–42	Sandy loam	
J. Thomas Way [4]	England	1850 [5]		
Greaves (*223, 330*)	Lee Bridge, England	1851	Mixture of loamy soil with sand and gravel	
G. von Mullendorf (*330, v. 11*).	Goerlitz, Germany	1853	Clay, loam, sand	Residual
E. Pfaff (*330*)	Erlangen, Germany	1867	Poor sandy soil	
E. Ebermayer [6] (*325; 330, v. 13, pp. 1–15*).	Munich, Germany	1868	Gravelly sand, fine sand, loessal loam, limestone sand, muck, humous garden soil.	
J. N. Woldrich (*330*)	Salzburg, Austria	Before 1870	Sandy loam, loam, sand.	
Do (330)	Oberdöbling, Austria	Before 1870	do	
J. B. Lawes and J. H. Gilbert (early work) N. H. J. Miller (later work) (*222, 223, 242, 252*).	Rothamsted, England	1870		Glacial, 8-inch cultivated soil over 10-inch friable clay over subsoil of rather stiff clay.
E. L. Sturtevant (*77, 84, 111*).	Waushakum Farm near South Framingham, Mass.	1875		Glacial
Levi Stockbridge (*82, 83*).	Amherst, Mass	1876–77		do
B. Latham (*247*)	Croydon, England	1878	Gravelly	Chalk soil, upland, alluvial.
E. Wollny (*330*)	Munich, Germany	1880	Sand, clay loam, humous sandy loam, mixture of sand and peat.	
Klucharov (*82*)	Moscow, Soviet Union			
A. Petermann (*401*)	Gembloux, Belgium	1881	Sandy loam	
Sturtevant and Babcock (*93, 122*).	Geneva, N. Y	1882	Dark clay loam	Glacial
Babcock and Goff (*93, 99*).	do	1887		do
Do	do	1887		do
M. P. E. Berthelot (*404*).	France	1887 [5]		
P. P. Dehérain (*405*)	Grignon, France	1888		
J. W. Sanborn (*85*)	Missouri Agricultural Experiment Station, Columbia, Mo.	1888	Clay loam	
B. Tacke et al. (*347*)	Ottersburg, Germany	1894	Peat	Peat
J. Hanamann (*346, 349*)	Lobositz, Austria	1896		

See footnotes at end of table.

installations 1688–1939

Lysimeter construction								
Type	Shape of surface	Dimensions				Material		Foreign substance in bottom layer
		Area		Depth		Walls	Bottom	
		Square meters	*Square feet*	*Meters*	*Feet*			
Filled-in	Round			0.19 .38 2.43	0.62 1.25 7.97	Lead		
do	do	0.051	0.545	.91	3.00	Tinned iron	Tinned iron	Gravel and sand.
do								
do								
do	Round	.165	1.77	.91	3.00	Cast iron		
do	do	.073	.79	.91	3.00			
do	do	.0003	.003	.38 .46	1.25 1.50			
do	Square	.836	9.0	.91	3.00	Slate		
do	do	.093	1.0	1.30	4.25			
do	Round	.018	.20	.15 .30 .61 1.22	.50 1.00 2.00 4.00	Tin		
do	Square	4.0	43.0	1.00 1.20	3.28 3.94	Concrete	Concrete, funnel-shaped.	
do	Round	.018	.20	.15 .30 .61 1.22	.50 1.00 2.00 4.00	Tin		
do	do	.018	.20	.15 .30 .61 1.22	.50 1.00 2.00 4.00	do		
Soil block	Rectangular	4.05	43.56	.51 1.02 1.52	1.67 3.34 5.00	4½-inch brick coated with cement.	Cast-iron plates perforated.	None.
do	do	20.234	218.0	.67	2.18			Do.
Filled-in		1.35	14.50	.91	3.00			
do	Square	0.837	9.0					
do	do	.04 .1	.43 1.08	.30 .50	0.98 1.64	Zinc	Perforated zinc.	
	Round			.20	.66	Metal cylinders.		
Filled-in	Square	1.0	10.76	1.00	3.28	Masonry		Gravel.
Soil block	do	.404	4.34	.91	3.00	Oak plank lined with heavy copper.	Sheet metal and copper.	None.
do	Round	.404 .404	4.34 4.34	.91 1.83	3.00 6.00	White-oak staves.	White-oak plank with copper.	
Filled-in	do	.404 .404	4.34 4.34	.91 1.83	3.00 6.00	do	do	
do		.152	1.64			Glazed earthenware pot.		
do		4.0	43.0	1.00	3.28	Concrete		
Soil block	Rectangular	40.46	435.6	1.30	4.26	Tarred fence	None except for drain tiles 7 feet apart.	None.
Filled-in	Round	.049	.529	.35	1.15	Glass	Glass	Broken glass.
do		.1	1.08	.50 1.00	1.64 3.28	Metal container.		

TABLE 2.—*Lysimeter*

Investigator and citation number in bibliography, or footnote	Location	Year of installation	Soil	
			Texture	Other characteristics
B. Welbel (*463, 474, 476*)	Ploti, Soviet Union	1899		Chernozem
Do	do	1903		do
Burt and Leather (*299*)	Cawnpore, India	1903	Loam, sand	Alluvium of Ganges River.
R. E. Horton (*103*)	Grafenburg, N. Y	1904	Clay, loam, sand	Clay from residual shale.
B. Welbel (*463, 474, 476*).	Ploti, Soviet Union	1905		Chernozem
Arnott and Leather (*284, 290*).	Pusa, India	1906	Loam, sand	Alluvium; contains 40 percent chalk.
Von Seelhorst (*351*)	Goettingen, Germany	1906 [5]	do	
Gerlach (*359, 378, 386*)	Bromberg, Germany	1906	Peat, loamy sand, sandy loam, humous sand.	
E. Krüger (*357-358*)	do	1908	Sandy soil	35 cm. surface soil, 85 cm. subsoil.
T. L. Lyon and J. A. Bissell (*109, 133*).	Ithaca, N. Y	1910	Silty clay loam	Dunkirk,[7] glacial
			Silt loam	Volusia,[7] glacial
			Sandy loam	Petoskey,[7] glacial
G. S. Fraps (*194*)	College Station, Tex	1910	Sand	Norfolk,[8] coastal plain.
			Fine sandy loam	Orangeburg,[8] coastal plain.
			Clay	Houston,[8] coastal plain.
			Loam	do
			Clay	Yazoo, bottom land
			Fine sandy loam	Miller (flood plain)
			Clay	Crawford [8]
			Fine sandy loam	Lufkin [8]
S. Collison and Seth Walker (*31*).	Gainesville, Fla	1910	Sand	Norfolk,[8] coastal plain.
C. A. Mooers and W. H. MacIntire (*152, 173*).	Knoxville, Tenn	1911	Loam	Hagerstown,[7] Limestone residual.
			Clay loam	Cumberland
			Silt loam	Memphis [7]
S. S. Peck (*55*)	Honolulu, H. I	1911 [5]		Upland
				Lowland
Von Feilitzen, Lugner, and Hjerstedt (*19, 456*).	Jonkoping, Sweden	1912	Muck	
J. Hendrick and H. D. Welsh (*263, 267, 271, 272, 276, 278*).	Aberdeen, Scotland	1914	Light	Glacial (granitic)
R. C. Collison, et al. (*108, 122, 125, 128*).	Geneva, N. Y	1914	Loam	Ontario [7] glacial
C. A. Mooers and W. H. MacIntire (*152*).	Knoxville, Tenn	1914	Same as 1911	

See footnotes at end of table.

installations 1688–1939—Continued

Lysimeter construction								
Type	Shape of surface	Dimensions				Material		Foreign substance in bottom layer
		Area		Depth		Walls	Bottom	
		Square meters	*Square feet*	*Meters*	*Feet*			
Filled-in	Square	.05	.538	0.30 .45 1.00	0.98 1.48 3.28	Zinc		
Ebermayer		.1	1.08	.25 .50 .75 1.00	.82 1.64 2.46 3.28	None		
Soil block		4.05	43.56	.91 1.83	3.00 6.00			None.
do	Round	.164	1.767	.91	3.00	Galvanized iron.	Galvanized iron on wooden rack	Do.
do				0.20 .25 .40 .50 1.00	0.66 .82 1.31 1.64 3.28			
do	Rectangular	4.05	43.56	.91 1.83	3.00 6.00	Masonry plastered with cement.	Galvanized sheetmetal.	None.
Filled-in		1.0	10.76	1.25	4.10	Iron	Iron	
do	Square	4.0	43.0	1.10	3.61	Masonry lined with Dutch tile.	Galvanized iron with 0.5-cm. holes.	None.
do	do	1.0	10.76	1.37	4.49	Iron sheets with reinforcements. Protected with asphalt tar.		10 cm. gravel.
do	do	1.619	17.4	1.30	4.25	Asphalt over concrete.		Sand and gravel.
do	Round	.073	.79	.61	2.00	Galvanized iron.	Galvanized iron.	
do	do	2.024	21.78	1.22	4.00	Steel		
do	do	.810	8.71	.31 .61 1.22 1.83	1.00 2.00 4.00 6.00	Heavy galvanized iron parafined inside.	Block tin outlet.	1-inch sand over glass wool.
do	do	.032	.35	.61	2.00	Galvanized iron painted with asphalt.	Perforated galvanized iron.	3-inch washed black lava sand.
do	Square	.64	6.87	.50	1.64			
Soil block	Rectangular	4.05	43.56	.99	3.25	Slate slabs	Perforated slate.	None.
Filled-in	Round	1.619	17.4	.61 1.22 2.44	2.00 4.00 8.00	Steel tanks lined with pitch.	Perforated metal disk.	Quartz over glass wool.
do	do	.202	2.17	.30 .61	1.00 2.00	14-gage galvanized iron painted with asphalt.		1-inch layer sand over glass wool.

TABLE 2.—*Lysimeter*

Investigator and citation number in bibliography, or footnote	Location	Year of installation	Soil	
			Texture	Other characteristics
H. K. Dean (*145*)	Umtilla, Oreg	1915	Medium sand, fine sand, coarse sand, silt, silt loam.	
E. E. Deturk [9]	Urbana, Ill	1916		
Miller and Duley (*86*)	Columbia, Mo	1918		
W. Geilmann (371)	University Goettingen, Germany.	Before 1920	Mild loam, yellow sand.	
C. A. Mooers and W. H. MacIntire (*158*).	Knoxville, Tenn	1921 [5]		
V. V. Gemmerling (*481*).	Moscow, Soviet Union	1922 [5]		Podzol
E. E. Deturk [9]	Urbana, Ill	1922–23		
W. B. Ellet and H. H. Hill (*198, 203*).	Blacksburg, Va	1922		Limestone soil
Do	do	1922	Silt loam	Frederick [7]
		1924	Sand	Hagerstown series [7]; Norfolk series [8]
J. E. Weaver and J. W. Crist.[10]	Lincoln, Nebr	1923	Fine sandy loam	
F. L. Duley (*87*)	Columbia, Mo	1925 [5]		
E. Krüger [9]	Berlin-Dahlem, Germany.	1925	Sandy	Glacial
M. Winnek (*308*)	Mikveli-Israel Experiment Station Palestine.	1925	Clay, loamy sand	Alluvial, glacial
H. H. Nicholson and B. Pantin (*256*).	Reading, England	1926	Loam	
H. J. Harper [11]	Stillwater, Okla	1927		
W. N. C. Belgrave (*309*).	Kuala, Lumpur, Federated Malay States.	1927	Sandy	Quartzite, eroded; laterite, good crumb structure.
A. W. R. Joachim (*301*, 1928: 3).	Peradeniya, Ceylon	1927		
W. C. Lowdermilk (*4*)	Berkeley, Calif	1927	Sandy loam	Holland
			Sandy clay loam	Aiken
			Clay loam	Altamont
V. P. Popov (*482*)	Soviet Union	1928 [5]		
H. C. Hendericksen (*147*).	Mayaguez, Porto Rico.	1928	Sandy clay	Practically devoid of organic matter and nutrients.
Niklas and W. Schropp (*382*).	Weihenstephan, Germany.	1928 [5]		
C. Pfaff (*398, 400*)	Limburgerhof, Germany.	1928	Alkaline, humous sandy soil, alkaline loam, acid slightly humous sandy soil, acid sandy loam.	
California Forest and Range Experiment Station.[13]	Devil Canyon Branch San Bernardino, Calif.	1929	Gravelly loam	Of granitic origin
F. Shreve (*2*)	Tucson, Ariz	1929	Adobe clay	Alluvial
M. F. Morgan (*14, 21*)	Windsor, Conn	1929	Loamy sand	Merrimac,[7] glacial
			Sandy loam	do
			Loam	Wethersfeld,[7] glacial
			Very fine sandy loam	Enfield,[7] glacial

See footnotes at end of table.

installations 1688–1939—Continued

Lysimeter construction								
Type	Shape of surface	Dimensions				Material		Foreign substance in bottom layer
		Area		Depth		Walls	Bottom	
		Square meters	*Square feet*	*Meters*	*Feet*			
Filled-in	Square	1.03	11.1	1.83	6.00	Concrete	Concrete	
do	do	25.29	272.25	2.51	8.25	do	do	
Soil block				1.22	4.00			
Filled-in	Round			1.30	4.26	Galvanized iron.		
do	do	.073	.785	1.83 2.14	6.00 7.00	Galvanized ingot iron tanks.	Conical shaped at bottom.	Coarse sand.
Ebermayer						None		
Filled-in	Rectangular	3.55	38.2	1.73	5.67	Concrete lined with asphalt.	Concrete	One-fourth inch crushed quartz.
do				.20 to .71	.67 to 2.33	Iron		Quartz pebbles.
do		.245	2.64	.20 to .71	.67 to 2.33			
Soil block	Round	.093	1.0	.91	3.00	Heavy galvanized iron cylinder.		None.
do	do	.186	2.0	1.22	4.00	Asphalt between soil and cylinder.	Perforated copper plate.	Lead shot.
Filled-in		1.0	10.76	1.50	4.92	Masonry	Masonry	
do								
Soil block								
Filled-in		0.0506	0.545	0.32	1.05	Glazed earthenware.		1-inch coarse washed gravel.
do								
Soil block								
Ebermayer						None		
Filled-in	Round	.067	.721	.43	1.41	Glazed earthenware.		
do	Rectangular	.929	10.00	.76	2.50	Galvanized iron.	Galvanized iron.	Gravel and sand.
Ebermayer						None		
Filled-in	Round	(12)				Iron	Iron	
do								
do	Square	1.0	10.76	1.0	3.28	Concrete	Concrete	10 cm. gravel, 50 cm. sand.
	Circular	.45	4.91	1.14	3.75	Galvanized iron.	Galvanized iron.	
Soil Block	Round	.067	.72	.43	1.41			
Filled-in	do	.206	2.22	.23 .51 .76	.75 1.67 2.50	Metal cylinders.		Pure quartz sand.
do	do							
do	do							
do	do							

TABLE 2.—*Lysimeter*

Investigator and citation number in bibliography, or footnote.	Location	Year of installation	Soil	
			Texture	Other characteristics
J. S. Joffe (*88, 89, 90, 91*)	New Brunswick, N. J.	1929	Sandy loam	Sassafras [7]
J. Bartels and W. Friedrich (*396, 397*).	Eberswalde, Germany	1929	Yellow fine sand	Glacial
California Forest and Range Experiment Station.[13]	Berkeley, Calif.	1929	Sandy clay loam Clay loam	Aiken Altamont
University of Pretoria (*489*).	Pretoria, South Africa	1930	Sandy loam	
A. Demolon and E. Bastisse (*419, 425, 427, 428*).	Versailles, France	1930	Loam	Upland
Arizona Agricultural Experiment Station (*1*).	Tucson, Ariz.	1930	Clay	Gila (alluvial) Mohave (pedocal)
G. W. Musgrave (*65*)	Temple, Tex.	1931		
California Forest and Range Experiment Station.[13]	Berkeley, Calif.	1931	Sandy clay loam Clay loam	Sierra Altamont
R. C. Collison (*131*)	Geneva, N. Y.	1932	Loam	Ontario,[7] glacial
H. A. Lunt (*19, 23*)	New Haven, Conn.	1932	Gravelly fine sandy loam.	Hartford, glacial
California Forest and Range Experiment Station.[13]	Berkeley, Calif.	1932	Clay loam	Altamont
Southwestern Forest and Range Experiment Station.[13]	Sierra Ancha Experimental Forest, Globe, Ariz.	1932	Sandy clay do Clay loam	Quartzite Granite Basalt
California Forest and Range Experiment Station.[13]	North Fork Branch Station, North Fork, Calif.	1932	Sandy clay loam	Sierra
G. W. Musgrave (*65, 69*).	Clarinda, Iowa	1933	Silt loam	Marshall, prairie soils Shelby, prairie soils
P. E. Karraker and C. E. Bortner (*71*).	Lexington, Ky.	1933	do	Maury [8]
J. E. Adams (*148*)	South Carolina Experiment Station, Clemson College, S. C.	1933		
California Forest and Range Experiment Station.[13]	North Fork Branch Station, North Fork, Calif.	1933	Sandy clay loam	Sierra
Do.[13]	San Dimas Experimental Forest, Glendora, Calif.	1934	Sandy loam	Holland (Pacific coast)
G. W. Musgrave [9]	Statesville, N. C.	1934		Cecil, Red soil of Piedmont.
F. Boeuf and V. Novikoff (*486*).	Ariana, Tunis	1934	Clay	Calcareous
E. Miege (*487*)	Rabat, Morocco	1934	4 soil types	
University of Pretoria (*489*).	Pretoria, South Africa	1934		
Intermountain Forest and Range Experiment Station.[13]	Boise Basin Experimental Forest, Idaho city, Idaho.	1934	Sandy	Granitic
Lake States Forest Experiment Station (*205*).	Upper Mississippi Valley Soil and Water Conservation Experiment Station La Crosse, Wis.	1934	Silt loam	Clinton

See footnotes at end of table.

installations 1688–1939—Continued

Lysimeter construction								
Type	Shape of surface	Dimensions				Material		Foreign substance in bottom layer
		Area		Depth		Walls	Bottom	
		Square meters	*Square feet*	*Meters*	*Feet*			
Ebermayer	Round	0.073	0.79	0.18 .42 .58 .80	0.6 1.4 1.9 2.6	None	Funnel	Quartz pebbles.
Filled-in	Square	1.0	10.76	1.50	4.9	Iron	Iron	10 cm. gravel.
do	Rectangular	1.858	20.00	.91	3.00	Redwood	Redwood	Gravel and sand.
do	Round	2.63	28.27	1.22 1.68	4.0 5.5	Pitch and tar mixture over cement plaster.		
do	Square	1.0	10.76	.60	2.0			
do	do	1.49	16.0	1.53	5.00	Concrete	Concrete	
Soil block	Round	.648	7.0	.91	3.00			Coarse gravel.
Filled-in	Rectangular	7.432	80.00	1.27	4.17	Iron (painted).	Iron (painted) on concrete foundation.	Gravel and sand.
Ebermayer	Round	.073	.79	.30 .58 .89 1.19	1.0 1.9 2.9 3.9	None	Funnel	
Filled-in	Round	.206	2.22	.10	.33	Galvanized iron.		
Ebermayer	Square	.103	1.11	.50	1.64	None	Square funnel.	
Soil block	Circular	.650	7.0	1.27	4.17	Galvanized iron.	Galvanized iron.	None.
do		.322	3.47	.56	1.83	Steel		Do.
Filled-in	Rectangular	.646	6.95	.53	1.75	Galvanized iron.	Galvanized iron.	Fine gravel and sand.
Soil block	Round	.648	7.00	.91	3.00	Metal		Coarse gravel.
Filled-in	do	.256	2.761	.61	2.00	Steel		
do	do	2.023	21.8	1.22	4.00	do		
do	Circular	.455 .646	4.90 6.95	.91	3.00	Galvanized iron.	Galvanized iron.	Fine gravel and sand.
do	Rectangular	20.234	217.8	1.83	6.00	Concrete	Concrete	None.
Soil block	Round			1.22	4.00	Metal		
Filled-in	Square	4.0	43.0	2.0	6.56	Reinforced concrete.	Concrete	
do								
Ebermayer	Round	0.552	5.94	1.22	4.00	None	Copper funnels.	
Filled-in	Rectangular	8.093	87.12	1.22	4.00	3/16-inch steel plate.	Concrete	Gravel.
Soil block	do	18.58	200.00	1.22	4.00	Asphalt-lined concrete.		Sand.

TABLE 2.—*Lysimeter*

Investigator and citation number in bibliography, or footnote	Location	Year of installation	Soil	
			Texture	Other characteristics
Southwestern Forest and Range Experiment Station.[13]	Sierra Ancha Experimental Forest, Globe, Ariz.	1934	Clay	
Do.[13]	do	1934	Stony sandy clay	Quartzite
Do.[13]	Fort Valley Branch Station, Fort Valley, Ariz.	1935	Clay loam	Basalt
S. Osugi (*313*)	Kyoto, Japan	1935[5]	Sandy loam, clay soil with gravel.	
R. S. Stauffer and R. S. Smith (*59*).	Urbana, Ill	1935	8 silt loams	Prairie soils
North Appalachian Experimental Watershed, Soil Conservation Service (*138, 139, 140*).	Coshocton, Ohio	1936	Silt loam	Muskingum,[7] Residual from sandstone and shale.
Alabama Experiment Station.[9]	Auburn, Ala	1936	Sandy loam, fine sandy loam, clay loam.	Norfolk[8], Hartsells[8], Decatur[8]
Southwestern Forest and Range Experiment Station.[13]	Sierra Ancha Experimental Forest, Globe, Ariz.	1937	Sandy clay loam	
Rocky Mountain Forest and Range Experiment Station.[13]	Manitou Experimental Forest, Colorado Springs, Colo.	1937	Gravelly	Granitic
California Forest and Range Experiment Station.[13]	San Dimas Experimental Forest, Glendora, Calif.	1937	Sandy loam	Holland (Pacific coast).
Do.[13]	do	1937	do	do
Do.[13]	do	1937	do	do
Do.[13]	do	1938	do	do
Do.[13]	do	1938	do	do
North Appalachian Experimental Watershed, Soil Conservation Service.[14]	Coshocton, Ohio	1938	Silt loam	Subsoils used: Loam, silt loam, silty clay.
Southwestern Forest and Range Experiment Station.[13]	Sierra Ancha Experimental Forest, Globe, Ariz.	1938		
Do.[13]	do	1939	Sandy loam	

[1] DE LA HIRE, PHILIPPE. See footnote 4, p. 9.
[2] MAURICE. See footnote 5, p. 9.
[3] GASPARIN. In Cours d'Agriculture, v. 2, p. 116. No date. [Original not seen.]
[4] WAY, J. T. ON THE POWER OF SOILS TO ABSORB MANURE. Roy. Agr. Soc. England Jour. 11: 313–379. 1850.
[5] Year published.
[6] EBERMAYER, ERNST. UNTERSUCHUNGS-ERGEBNISSE ÜBER DIE MENGE UND VERTEILUNG DER NIEDERSCHLÄGE IN DEN WÄLDERN. Forstl. Naturw. Ztschr. 6: 283–301. 1897.

installations 1688–1939—Continued

Type	Shape of surface	Lysimeter construction						Foreign substance in bottom layer
		Dimensions				Material		
		Area		Depth		Walls	Bottom	
		Square meters	*Square feet*	*Meters*	*Feet*			
Filled-in	Rectangular	9.29	100.00	2.44	8.00	Concrete	Concrete	Sand.
Soil block	do	83.61	900.00	.91 to 1.83	3 to 6	do		None.
do	do	.312	3.47	.56	1.83	Steel		Do.
do	do	.83	8.9					Do.
do	Round	.648	7.0	1.02	3.3	Galvanized iron.		Crushed quartz.
do	Rectangular	8.089	87.08	2.44	8.00	Concrete lined with asphalt.	Perforated steel plate.	None.
Filled-in	Round	.455	4.9	.30 .91	1.00 3.00	Galvanized iron painted with asphalt.	Galvanized iron.	Gravel and sand.
Soil block		.646	6.95	.71	2.33	Steel		None.
Filled-in	Rectangular	.405	4.36	.46 .91	1.50 3.00	do	Steel	Gravel.
do	Circular	.484	5.21	1.01	3.33	Galvanized iron.	Galvanized iron.	None.
do	do	.161	1.73	.61	2.00	do	do	Do.
Filled-in, unconfined.	Rectangular	28.451	306.25	2.13	7.00	Soil		Do.
Filled-in	Rectangular	2.230	24.00	1.83	6.00	Concrete	Concrete	Do.
do	Circular	.161	1.73	.61	2.00	Galvanized iron.	Galvanized iron.	Do.
do	Round	.032	.349	1.04	3.42	Glazed tile	Wire screen over sheet metal.	Washed sand.
Soil block		.646	6.95	.71	2.33	Steel	Steel	None.
Filled-in		41.80	450.00	1.22	4.00	Concrete	Concrete	Sand.

[7] Gray-Brown Podzolic soils.
[8] "Red and yellow" soils.
[9] Private communication.
[10] Weaver, J. E., and Crist, J. W. See footnote 4, table 1.
[11] Harper, H. J. See footnote 6, table 1.
[12] 30-gallon capacity.
[13] Unpublished data through courtesy of U. S. Forest Service.
[14] Unpublished data.

TABLE 3.—*Chronology of the development of lysimeters, 1688–1937*

Type of lysimeter	Investigator	Location	Year of construction
First lysimeter	De la Hire	Paris, France	1688
First lysimeter with run-off provision	Dalton	Manchester, England	1796
First extensive use of lysimeters for the study of soil chemistry.	Way	England	1850
First soil-block lysimeter	Lawes and Gilbert	Rothamsted, England	1870
First (soil-block) lysimeter in the United States.	Sturtevant	South Framingham, Mass.	1875
First large comparative lysimeter study	Wollny	Munich, Germany	1880
First lysimeter in natural contact with substrata drained by tiles.	Sanborn	Columbia, Mo	1888
First Ebermayer lysimeter	Welbel	Ploti, Soviet Union	1903
First soil-block lysimeter with run-off provision.	Leather	Cawnpore, India	1903
First lysimeter, with weighing mechanism	V. Seelhorst	Goettingen, Germany	1906
First soil-block lysimeter with weighing mechanism.	Weaver and Crist	Lincoln, Nebr	1923
First Ebermayer lysimeter in the United States.	Joffe	New Brunswick, N. J	1929
First soil-block lysimeter with self-recording weighing mechanism.	Research Division, Soil Conservation Service.	Coshocton, Ohio	1936
First lysimeter with tensiometers for measuring capillary tension of soil moisture.	Iowa State College and Soil Conservation Service.	Clarinda, Iowa	1937

During the nineteenth century lysimeter investigations became more frequent. All investigations conducted before 1870 were with filled-in lysimeters. This is what should be expected, since pedology was in its infancy; and, in general, only the surface soil was subjected to research by the pioneers of soil science. The plow layer is disturbed regularly through cultivation, and therefore it seemed unobjectionable to remove it from its original site and place it in lysimeters. At that time the significance of the soil profile as a nonseparable unit was not understood. De la Hire, Dalton, and Maurice, as well as a number of other investigators in England, Germany, Austria, and France, used lysimeters of the filled-in type merely for the study of the rate and amount of the percolate. They did not pay much attention to the chemical composition of the percolate, and most of them did not seem to care greatly whether the soil in the lysimeters was covered with vegetation or not.

In 1850 Way,[6] in England, reported his fundamental studies on the chemical changes of solutions percolating through soil. He used a number of laboratory lysimeters, filled them with different types of soils, and found that bases of the sprinkling solution were exchanged against the calcium of the soil and that the power of soils to absorb fertilizing elements depends on its clay content. He also noticed that burning destroyed this capacity of the clay. Through this work, Way laid the foundation for further base-exchange studies. He was the first to carry out extensive lysimeter investigations to clarify the problem of soil fertility. The influence of his studies is noticeable in the work begun 20 years later at Rothamsted.

Lysimeter investigations took a great step forward in 1870, when Lawes, Gilbert, and Warington established three "drain-gauges" at Rothamsted, England. These men recognized that basic information concerning the natural rate and chemical composition of drainage water could be obtained only from natural, undisturbed soils. In-

[6] See footnote 4, table 2.

stead of filling soil into tanks they left the soil intact and placed a perforated iron bottom under it and masonry walls around it. These lysimeters exceeded all preceding installations in area five times, and they are responsible for a great many findings concerning the percolation rate of rain water and the leaching of plant nutrients from soil. After 68 years of successful operations, these lysimeters are still in use. It is not surprising that the construction of these first soil-block lysimeters met with some difficulties and that leakage through the side walls could be stopped only after repeated repairs. As in all early installations, run-off is not allowed to occur from the Rothamsted lysimeters. Under English climatic conditions, this may not be a great handicap, since the rainfall is generally much more gentle than in this country. The depth of each of these "drain gauges" is different; therefore, no replicate exists to check directly the results obtained. However, a comparison of the data from these lysimeters with those from later installations indicates a high degree of reliability of the Rothamsted work. While the construction of these lysimeters was very satisfactory, the fact that only three of them exist and that these three are located on the same soil type and are otherwise under identical conditions, makes it impossible to interpret the effect of soil type, vegetation, and various other factors on the rate and amount of percolation.

Research to investigate such effects was started by Wollny (*330*) in Munich, Germany, in 1880. In his very elaborate set-up he studied the following problems:

(1) Influence of soil structure on the rate and amount of percolate.
(2) Effect of soil type on the rate and amount of percolate. (Sand, clay loam, humous sandy loam, peat, a mixture of sand and peat, stony sand, stony humus and sandy loam were used.)
(3) Effect of soil depth on the rate and amount of percolate.
(4) Effect of soil surface cover on the rate and amount of percolate. (Sand, loam, dead vegetable mulch, and growing vegetation were compared.)
(5) Effect of the distribution of precipitation on the amount and rate of percolate and on the ratio of percolate to precipitation.

This study, continued through almost a decade, was carried out with a large number of lysimeters of the filled-in type. Wollny was a very thorough worker who studied many questions of agricultural meteorology and soil physics, and it is not surprising that the majority of his lysimeter findings have been verified by later workers. The great number of phenomena that he subjected to research excluded for him the use of soil-block lysimeters, which would have been altogether too costly. The use of the filled-in type cannot be considered very objectionable for the study he carried out, since he wanted to obtain comparative information concerning natural percolation. It is to be noted that Wollny did not include any analyses of the leachate in his research.

For many purposes the filled-in type of lysimeters proved to be inadequate, as, for instance, the study of the moisture relationships in forests. Ebermayer [7] (*325, 330*) carried out such studies in numerous meteorologic stations in the forests of Bavaria. He found that filled-in lysimeters could not depict natural conditions, since the tree roots did not reach into the lysimeters and hence could not remove water from the soil. Within a few seasons the lysimeter soil became

[7] See footnote 6, table 2.

so waterlogged that the experiment had to be discontinued. This may have been due not only to the lack of tree roots, but also to the poor under-drainage inherent in most lysimeters. This phenomenon is discussed in detail on pages 19–22. In order to overcome the obstacles encountered in the study of hydrologic conditions of forests, Ebermayer suggested a rather unique scheme. A large area of the forest, about one-fourth hectare, was to be undermined about 1.5 meters below the soil surface so that the soil and trees would remain undisturbed. The roof of this wide tunnel would consist of impervious material and would slant to one point where the percolate could be measured. This project evidently did not materialize, owing to the technical and financial difficulties involved in such a huge undertaking.

A number of years later, Russian investigators modified this plan of collecting percolate from undisturbed, unencased soil. In particular, the size of the collecting funnel was considerably reduced and thus construction problems were simplified. From Russia (*463*, *474*, *476*, *481*, *482*) this funnel-type lysimeter came to the United States (*88*, *89*,), and it has, therefore, frequently been called a Russian lysimeter, although a Russian investigator, Katchinsky,[8] was among the first to call it the Ebermayer type.

The first lysimeter in the United States was built by Sturtevant on his farm in Massachusetts in 1875, and it is interesting to note that this was of the soil-block type. In 1882 Sturtevant installed another set of three lysimeters at the experiment station at Geneva, N. Y. These lysimeters were also of the soil-block type and were transported from the original location to the lysimeter pit. They were constructed of white oak planks lined with heavy sheet copper and had a bottom made of a heavy flat section of "boiler iron." After a number of years, Babcock and Goff, who continued this work, believed that the lysimeters dried out too readily, and they therefore built four additional lysimeters with an artificially maintained ground-water table. Two of these lysimeters were of the soil-block type and two were filled with soil that was previously sifted and dried. The records obtained from this research do not seem to have been very convincing; for the installation was abandoned in 1890.

None of the earlier lysimeters was provided with run-off facilities, except, of course, the Ebermayer lysimeters, whose lack of side walls allows free run-off. It is significant that the first large lysimeter with run-off provision was constructed in India (*284*, *290*), where dry periods alternate with seasons of intense storms. This installation was made in 1903 and has been followed by numerous others, particularly in the United States, where intense rainfall frequently causes large quantities and high rates of run-off.

Direct information can be obtained on only percolation and run-off from stationary lysimeters. The sum of evaporation and transpiration can be calculated for longer periods by differences if precipitation data are available, but accurate evapotranspirational data for short-time intervals can be obtained only if the lysimeters are weighed, as only then will the change of soil-moisture content be known. Von Seelhorst (*351*) was the first to equip lysimeters with weighing devices (1906). Other German investigators (*357*, *396*, *397*) [9] used similar

[8] KATCHINSKY, N. A. METHODES POUR DETERMINER LA PERMÉABILITÉ DU SOL À L'EAU EN VUE D'UNE IRRIGATION. 1st Comn. Internatl. Soc. Soil Sci. Trans., section Soviet, Moscow, A, 2: 79–99. 1934.

[9] See also footnote 3, p. 4.

installations for the study of transpirational losses of various crops and the evaporation from bare soil. Weaver and Crist [10] weighed soil-block lysimeters in this country (1923), and the first soil-block lysimeters with self-recording weighing mechanism were built in 1937 by the Soil Conservation Service at Coshocton, Ohio.

These large lysimeters (one five-hundredths of an acre) at Coshocton, in the planning of which the writers took part, embody numerous features recognized as essential for a study of true natural percolation and other hydrologic factors. In order to illustrate how some of these features have been incorporated, a short description of these lysimeters will be given. (Details of design and construction have been reported by other members of the Research Division, Soil Conservation Service (*138*, *139*, *140*). Figure 1 shows a schematic cross section through an entire lysimeter installation. The lysimeter is 8 feet deep; the soil extends merely 2 to 3 feet; disintegrated rock extends another 2 to 3 feet; and the lowest 3 to 4 feet contains undecomposed parent rock. All these layers are left in their natural structure, and the rock rests immediately on the percolate sieves. Water reaching these strata can fairly be called percolation, and relative assurance is given that no plant roots reach this depth. It seems probable that water passes with little difficulty from the rock fissures into the multiple percolate collectors. In figure 1 are shown cross sections of steel strips that serve to prevent fictitious percolation from descending in the space that may at times exist between the soil block and the side walls of the lysimeter.

Provision is made for the collection of run-off, and the amounts of water and soil in the run-off tank are recorded. Conditions with respect to run-off in the lysimeter are not representative of field conditions because the vertical side walls cut off overland flow. To date, however, no method that permits overland flow to occur on a lysimeter has been found to give an accurate knowledge of the amount of water that infiltrates into the soil mass. The side walls have another disadvantage, that of creating a border effect, which is due both to the lack of vegetation on a strip about 15 inches wide and also to cutting off the soil from its surroundings. The necessity of isolating a soil block for the study of its percolation and its weight changes requires such a procedure. An attempt is made to keep the temperature in the subsoil somewhat the same as it would be in nature by keeping the service tunnel closed as well as possible. Grease seals (fig. 1) are placed between the top of the lysimeter walls and the outside retaining wall to keep the atmospheric air from descending into the tunnel and at the same time to allow for free play of the lysimeter on the scales. These record weight changes accurately to the nearest 5 pounds, which is equivalent to approximately one one-hundredth of an inch of rainfall.

DISCUSSION

The purpose of lysimeter investigations is twofold; the measurement of the hydrologic balance of the soil, in which the amount and rate of percolation is the primary consideration, and the determination of the chemical losses accompanying percolation. Omitting laboratory lysimeters, which are used to study special percolation problems and

[10] See footnote 4, table 1.



FIGURE 1.—Schematic cross-sectional diagram. A soil-block lysimeter with self-recording weighing mechanism used at the North Appalachian Experimental Watershed, Coshocton, Ohio.

which are sometimes sprinkled artificially, lysimeters are set up to supply data that correspond closely to those that would be obtained under natural conditions. It is surprising to note that in the majority of the lysimeters soil and percolation conditions prevail that are far from natural, and to sanction the use of such lysimeters by declaring very shallow filled-in lysimeters without run-off provision to be standard (*210*) appears to be unjustified. The removal of soil from its original site, and the process of sifting, mixing, and replacing it in a container is bound to change the resulting soil climate considerably. If this mixing is restricted to the plow layer, it does not seem to be so objectionable, since it is this part of the soil that is likewise disturbed in the field by agricultural implements. The mixing of the plow soil in itself cannot adversely affect the results, but its separation from the underlying soil layers is apt to change the percolation rate. To overcome this objection, some investigators have filled lysimeter tanks with a number of soil horizons in the same sequence as they occur in the natural profile, but in this way it frequently happens that soil that was on the upper side of a certain horizon will be placed in the lower part of the same horizon when filled into the lysimeter tank. This will result in a change of chemical, physical, and biological conditions and is apt to influence the rate of percolation and its chemical composition.

The change of the original soil structure, the obviation of root channels and of soil fissures in the lower horizons are apt to decrease greatly the rate of percolation, particularly in heavy subsoils; but the changes of percolation from natural conditions are not restricted to filled-in lysimeters. The greatest functional error in all three types of lysimeters probably occurs at the boundary of the lysimeter soil and the air beneath it. When the gravitational water reaches this point it will have to overcome the resistance set up by surface tension before it can leave the soil. This phenomenon has been studied by various soil investigators and has been described particularly well by Lebedev [11] and by Zunker [12] (*389*), but it appears that many lysimeter installations have been planned without a complete comprehension of this drainage-impeding effect of the soil-air boundary at the bottom of the lysimeter.

Frequently sand or gravel is placed below the soil to keep it from washing out through the drain pipe. This practice is also followed in an attempt to provide for better drainage. The writers have investigated the effect on drainage by covering the bottom of lysimeters with 6-inch layers of washed sand and placing various layers of soil above them. At the end of the dry season (October) these layers of soil and sand were removed and the soil-moisture content of numerous samples determined gravimetrically. An accumulation of soil moisture was found above the soil-sand boundary as well as above each boundary between soil layers of different texture and a particularly great accumulation of water was present in the lower part of the sand just above the bottom screen.

In very humid climates where percolation rates are high the percolate restriction at the soil-air boundary will not be very serious,

[11] LEBEDEFF, A. F. THE MOVEMENT OF GROUND AND SOIL WATERS. 1st Internatl. Cong. Soil Sci., Washington, 1928. Proc. 1: 459–494. 1928.

[12] ZUNKER, F. DAS VERHALTEN DES WASSERS ZUM BODEN. Blank's Handbuch der Bodenlehre 6:125. 1930.

because enough water percolates to overcome the surface-tension impediment and because the natural soils are moist most of the time. In drier climates, however, the soils in lysimeters will have a tendency to contain much more moisture in the bottom layer than they contain in this layer in the undisturbed profile. Unless lysimeters are very deep, the roots of the plants growing on them will reach this excess water and hence will supply the plants with more water for transpiration and more nutrients than they would under natural conditions. In this way the rate and amount of percolation and the chemical composition of the percolate are changed. Frequently, this accumulation of water, together with the prevention of run-off, produces such an unhealthful condition that plant growth suffers, and the rate of transpiration is reduced below normal. Some investigators even aggravate this objectionable waterlogged condition of lysimeter soils by preventing the growth of higher plants and thus eliminating the natural disposal of water through transpiration. After a few seasons, moss and algae take the site, which has been made unsuitable for agricultural crops. It is a well-known fact that waterlogging of soils is accompanied by many changes, such as lowering of the oxidation-reduction potential, denitrification, dissolving of iron and titanium, formation of hydrogen sulfide, raising of the partial pressure of carbon dioxide, all of which will tend to alter the chemical composition of the percolate. Where side walls do not permit the horizontal flow of water in lysimeters, as in the monolith and filled-in types, the water will be forced to escape through the open bottom after sufficient hydrostatic head has been developed to overcome the drainage-impeding effect of surface tension. In Ebermayer lysimeters it may frequently happen that the water, when encountering the obstacle to percolation set up by the soil-air interface, follows the path of least resistance and flows around the funnel. This may explain the extremely low percolation rates reported by Joffe (*90*) in the lower soil horizons. Harper [13] found that under certain soil conditions Ebermayer lysimeters did not yield any percolate, whereas under other conditions the same lysimeters acted as drains and therefore produced excessively high percolation rates. It is doubtful whether Ebermayer lysimeters will yield reliable data for the study of the composition of the percolate, but it seems certain that they cannot yield quantitative hydrologic data.

While this impeding effect will be operative in a greater degree in dry seasons and climates, the large amounts of gravitational water that percolate in times of excessive precipitation or after the thawing out of the soil in the spring will find less resistance at the soil-air boundary than they would in passing downward through an almost saturated subsoil. After such excessive drainage has taken place, no ground water is at the disposition of the lysimeter soil; and this soil may dry out considerably, whereas natural soil would have been kept moist for a much longer period. It is these two opposing factors, the impediment of percolation by the surface tension operative at the soil-air interface and the lack of resistance to percolation below the lysimeter, that frequently give percolation data of apparently satisfactory agreement with natural conditions, when only the amount

[13] HARPER, HORACE J. A STUDY OF LYSIMETER INSTALLATIONS FOR THE MEASUREMENT OF GRAVITATIONAL WATER. Unpublished paper presented before the meeting of the Amer. Soc. Agron., November 1937.

of percolation of longer periods is considered. But data, obtained through compensation of two large errors, are of little value.

One of the greatest objections to lysimeter investigations is that, generally, measurements of the moisture content of the various soil layers are not made, and therefore no knowledge can be obtained of the movement of the water within the lysimeter and no comparisons made with the moisture content of similar soils in natural surroundings. Determinations of gravitational soil moisture are normally out of the question in a lysimeter. Recently, however, tensiometers have been used (*69*) to study soil-moisture conditions in lysimeters, and other work is under way on this problem.

It is probable that air-pressure conditions in most lysimeters are different from those in the undisturbed soil, since so many factors occur in lysimeters that are likely to change soil-air pressure. The most important of these factors are probably the open bottom and the side walls. That such pressure changes influence percolation rates cannot be doubted, but it is likely that their magnitude is much smaller than the changes due to the factors mentioned above. Some careful research will be necessary to clarify this problem.

Soils containing a fair percentage of clay shrink on drying. When they are contained in a lysimeter it frequently happens that fissures appear between the soil and the side walls, and precipitation often results in direct percolation along the side walls before the soil has had time to swell and fill these openings. The amount of such "fictitious" percolation can be very large, and precautions should be taken to prevent it.

Precipitation is not the only source of soil water; condensation and absorption of water vapor also contribute a significant share, and equipment should be installed to ascertain the magnitude of these accretions. If only chemical phenomena are investigated, such information is not particularly important. Wherever possible, provision should be made for weighing the lysimeter. Scales with weight recorders are to be preferred, since they permit the study of diurnal moisture fluctuations in the soil body. The following equations show how the data from a weighing lysimeter can be used.

Water addition	= Water loss	+ Water storage
Precipitation (direct measurement)	Run-off (direct measurement)	Direct measurements by weight [14]
Rain	Percolation (direct measurement)	
Snow	Evapo-transpiration [15]	
Hail	Evaporation	
Sleet	Transpiration	
Fog		
Condensation		
Dew (plants)		
Condensation (soil)		
Adsorption (soil)		

Precipitation + condensation = run-off + percolation + evapo-transpiration + storage [16]

Precipitation − run-off − percolation − storage = evapo-transpiration − condensation

[14] Weight changes of a lysimeter may be caused by other factors besides water. Most of these are insignificant, but such factors as application of manure, soil erosion, growth, and harvest of plants have to be included in the calculation.

[15] Evaporation and transpiration cannot be separated accurately in a single lysimeter experiment.

[16] Storage may be positive or negative (loss of weight). If it is negative, the opposite sign should be used.

These equations are applicable to any period of time studied. All factors on the left-hand side of equation (3) can be determined directly, while only the difference between evapo-transpiration and condensation can be obtained with this equation from the data of a weighing lysimeter. A rather close separation between condensation and evapo-transpiration can be obtained by plotting accumulated evapo-transpiration minus condensation against time, wherever accurate weights are recorded for 30-minute intervals or less.

ESTABLISHED LYSIMETER FINDINGS

A critical review of the literature on lysimeters reveals that some authors have studied phenomena that have been satisfactorily investigated by others at previous times. All the factors that depend on purely local conditions obviously have to be studied at each given location. This is true of the determination of the magnitude of data obtained through lysimeter research. But since many lysimeters are constructed that cannot possibly represent natural hydrologic conditions, their data will yield at best only generalizations, and of these a good many are already known. A summary of those fundamental facts that have been established through the lysimeter research of two and a half centuries is presented.

FINDINGS WITH RESPECT TO THE MOISTURE RELATIONSHIPS OF SOILS[17]

TEXTURE

Percolation is more rapid through light soils than through heavy soils. Percolation is more continuous through heavy soils than through light soils. The total amount of percolation per year is greater through light soils than through heavy soils.

Heavy soils lose more water through evaporation and run-off than light soils.

STRUCTURE

Porous crumb structure favors percolation, whereas single-grain structure tends to decrease percolation. Exchangeable cations affect the percolate indirectly as they affect structure and swelling. Calcium and hydrogen soils have a higher rate of percolation than the same soils when treated with sodium salts, because the deflocculation of the clay brought about by the sodium ion results in the elimination of many of the larger pore spaces.

OTHER SOIL CHARACTERISTICS

Natural fissures, worm holes, and channels of dead roots increase percolation. Virgin forest soils allow a much greater percolation to occur than the same soils after they have been cleared and cultivated.

The amount of drainage is largely a reflection of the amount of precipitation and of the total amount of dry matter removed in crops, being increased by increases in precipitation and decreased by increases in the amount of dry matter removed. The influence of the depth of the soil column on percolation is largely determined by soil type. Since various investigators have reported entirely different correlations of soil depth with percolation, no general statement can be made.

[17] The presentation of lysimeter findings is based on a comparative study of the data cited in this discussion of lysimeters.

Fertility affects the amount and rate of percolation only indirectly through the growth of plants. In general, high fertility is associated with increased transpiration and decreased percolation.

SOIL COVER

When a lysimeter soil is covered with coarse sand or gravel, infiltration is increased and evaporation decreased, and hence percolation is increased; this relationship might be obscured or changed by plant growth if the plants take up the greater amount of available water.

A mulch of dead vegetal material shows an effect very similar to a mulch of sand or gravel. The beneficial effect of a dead vegetal mulch on the moisture relationships of soils increases with increases in the thickness of the mulch to about 3 inches; but when the mulch gets much thicker the amount of water intercepted and directly evaporated surpasses the amount of water that is saved from evaporation, and hence percolation decreases.

Owing to transpirational losses, lysimeters covered with living vegetation usually show less percolation than those without vegetation. This difference is more or less proportional to the rate of growth, increasing during the summer and becoming negligible during the winter.

WEATHER AND CLIMATE

An annual cycle of percolation rates is frequently observed. In the Temperate Zone, percolation is greatest in the latter part of the winter and in the early spring and decreases rapidly with the growth of plants in the summer. In many installations percolation ceases entirely in midsummer and does not begin until late in the autumn.

The amount of annual percolation from any given lysimeter generally fluctuates approximately to the same extent as precipitation, whereas evapo-transpiration remains rather uniform under similar conditions of plant cover. The amount of percolation is affected after extremely dry or extremely wet years in proportion to the change of the soil-moisture storage from normal.

As is consistent with the genesis of soils, percolation occurs freely through Pedalfers, whereas through Pedocals it occurs only occasionally and in small quantities, unless the soils have been irrigated heavily.

FINDINGS WITH RESPECT TO THE CHEMICAL CONSTITUENTS OF THE PERCOLATE

The rate of loss of chemical constitutents from lysimeter soils is dependent on the amount of percolating water. During the summer, when the rate of percolation becomes very low, the rate of loss of chemical constitutents also drops considerably, but the total concentration normally increases with the decrease of the percolation rate. This, however, is not true of all individual elements. The amounts in which the various substances occur in the percolate depend on their abundance, their affinity to the soil, their solubility, the temperature, and the rate of percolation.

In general, the amount of chemical constituents lost is greater from bare soils than from soils supporting a vegetation, because the plant roots take up a considerable part of the dissolved chemicals.

170523°—40——4

The difference of concentration is particularly great in easily leachable materials, such as nitrate, chloride, and sulfate, whereas phosphate, usually in low concentration, is leached in about the same small quantities from fallow lysimeters as from vegetated lysimeters. The exchangeable bases take up an intermediate position in this respect.

EXCHANGEABLE BASES

In every lysimeter installation of which the writers have studied percolate analyses, calcium is leached out in much greater quantities than the other bases. Magnesium, potassium, and sodium usually follow in the order named. It is probable that in saline soils sodium would be lost to a greater extent than calcium, but lysimeter experiments on such soils are rather uncommon, probably because of the very nature of Pedocal soils, in which percolation is very low.

NITROGEN

Nitrogen occurs in the percolate mainly in the form of nitrate. A part of the ammoniacal nitrogen is utilized by plants, and the remainder is held largely in the exchange complex. Any ammonia released from these sources is normally oxidized before leaving the soil, and hence little ammonia appears in the percolate. In peat soils, because of their reducing power, a great deal of the nitrogen in the percolate appears in ammoniacal form. Under conditions favorable for denitrification, nitrates are found in the percolate. In filled-in lysimeters it is frequently noticed that in the first year or two the nitrate content of the percolate is rather high, but afterwards it decreases to very low rates. This decrease is due to aeration caused by stirring and mixing the soil before it has settled and to the larger pore spaces remaining in the soil, which tend to make conditions favorable for nitrification. Later on, these spaces close up; and since the natural fissures and cracks have been destroyed by the sifting and filling process, aeration conditions, and hence nitrification, become poorer, particularly if drainage in the soil is impeded, as it is in numerous lysimeters.

PHOSPHATE

Phosphate is found only in traces in the percolate from most lysimeters irrespective of the fertility status of the soil and regardless of whether plants are growing on it. Only in case of high application of phosphate to a very light soil is the phosphate content of the percolate somewhat higher.

SULFATE

Rain water in practically every locality contains appreciable quantities of sulfate, particularly in the vicinity of cities and industrial plants, and it seems that the amount of sulfate in the percolate is somewhat dependent on the amount of sulfur in the rain. Other important sources of sulfur are commercial fertilizer and manure. Sulfate occurs in the soil in various compounds, some of which are not easily soluble, but there is usually an abundance of bases with which it can form soluble compounds, and, consequently, percolation waters are almost always relatively rich in sulfate. The amount of sulfate percolation, as well as nitrate percolation, is greatly affected

by plant growth. The rate at which mineralization of organic matter takes place is another factor affecting the amounts of sulfate in the percolate.

CHLORINE

Chlorine is supplied to the soil mainly through precipitation and fertilizers, and since most chlorides are very readily soluble, they are leached out of the soil in about the same quantities in which they are supplied. If lysimeters are cropped, some chlorine is retained by the plants. This, however, does not greatly change the amount of chlorides in the percolate, since plants absorb very little of this ion. The concentration of chlorides in percolating waters of any given soil does not change very greatly during the season if no fertilizer containing chlorides is added. The rate at which chlorides are lost from the soil is almost proportional to the rate of percolation. In any one year the amount of chlorides in the percolate may be smaller or larger than the amount added to the soil through precipitation and fertilizers, depending on the amount of percolating water, which in turn depends largely on the rainfall.

SILICA

Only a few investigators have analyzed silica in lysimeter leachings, and little is known in this regard. Silica is lost in great quantities from most soils. Low concentrations of electrolytes in the soil solution seem to encourage the percolation of silica, because under such conditions the inorganic soil colloids become easily peptized and comparatively mobile, which results in a greater loss of silica in the winter than in the summer.

OTHER ELEMENTS OR COMPOUNDS

Iron, aluminum, titanium, manganese, carbonate, bicarbonate, and organic carbon have been determined in lysimeter leachates by a few investigators, but in the authors' opinion no general conclusions drawn from the data published to date are warranted.

REACTION OF THE PERCOLATE

In general, the pH value of the percolate from an unfertilized soil is higher than that of the soil itself. This is due probably to the presence of basic carbonate or bicarbonate salts in the percolate. When fertilizers are applied to the soil the pH value of the percolate may be either higher or lower than that of the soil, depending on the physiologic reaction of the fertilizer. The reaction of leachings from soils treated with sodium nitrate is near the neutral point, whereas leachings from soils treated with ammonium sulfate are distinctly acid.

SUGGESTIONS FOR FUTURE LYSIMETER STUDIES

CONSTRUCTION OF LYSIMETERS

Two and a half centuries of lysimeter studies have greatly helped to promote the knowledge of the phenomena of the movement of water through soil and the leaching of chemicals from soils. Not in every case, however, does the yield of information obtained from a lysimeter investigation seem to correspond to the effort put into it. This is not surprising if one considers the many difficulties encountered

in the construction and operation of lysimeters. In the following pages an attempt is made to enumerate the factors that are of primary importance in the construction of lysimeters.

In general, lysimeters should be constructed in such a way that their moisture relationships correspond closely to those of soils under natural conditions. (In some instances, special features that require a somewhat artificial environment have to be studied. The laboratory lysimeters will not be discussed here.) It should be considered essential that the soil in a lysimeter be in its original undisturbed condition. This is true regardless of whether the main purpose of the study relates to questions concerning hydrology or fertility. After the soil has once lost its identity by mixing, sifting, and packing, it is futile to hope that it will be restored to its original composition and structure by allowing it to settle for a few years. Flodkvist [18] found an increased permeability in soil that had been allowed to settle for 48 years. Nitrification is a very sensitive index of soil conditions. The percolate of freshly filled lysimeters, as has been mentioned, usually contains large quantities of nitrates, since the soil was aerated when it was handled and its pore space is greater than it was when the soil was in its original position. After a few years the yield of nitrates in the leachate rapidly decreases to a point below that which would be obtained from the natural soil. The soil has then become so packed and dense that aeration proceeds only with difficulty. The cracks that provide air circulation in a natural soil have not had time to form. The change in nitrification can be very readily observed. There is no doubt that most of the other physical, chemical, and biological soil reactions are seriously affected by filling the soil into lysimeter boxes. The heavier the soil and the more mature its profile the greater will be the disturbance created by digging and replacing, even if an attempt be made to keep the horizons in their original sequence. It can be assumed that a sandy soil, particularly if its profile is rather undeveloped, will be little affected by removal from its original site and replacement in a lysimeter. In general, however, a soil should be left unmolested and the lysimeter built around it if data are to be obtained that will serve to answer the intricate problems of the moisture relationships of natural soils.

It can hardly be doubted that the separation of the soil profile from the substrata and the placing of a perforated bottom under it will affect the rate of percolation. The reasons for this have been discussed. It would appear reasonable to follow the example of Sanborn (*85*), who in 1888 built large lysimeters merely by encasing one one-hundredths of an acre of natural soil in impermeable vertical walls and placing drain tiles at a 4-foot depth in the ground to collect the percolate. It seems that no other investigator has employed this type of lysimeter, probably because of the uncertainty of all the percolate being collected in the tiles and the probability of obtaining data that are not quantitative. The excessive drainage caused by the drain tiles can hardly be considered a valid argument against the Sanborn lysimeter, since some other lysimeters also are defective in this respect, at least during the wet season.

In the authors' opinion, it would be worth while to make a study of the usefulness of Sanborn lysimeters; for they permit a study of per-

[18] Flodkvist, H. AGRONOMISCH-HYDROTECHNISCHE ERGEBNISSE VON DRÄNUNGSVERSUCHEN AUF TONBODEN. 3d Internatl. Cong. Soil Sci., Oxford, Trans. 3: 164–168. 1936.

colate obtained from soil with a minimum of disturbance of the profile. Sanborn was not able to make such a study, because he left the Missouri Agricultural Experiment Station shortly after completing the lysimeters. As all other lysimeters are constructed with an artificial bottom and as percolation of water through soil cannot be studied at present without lysimeters, it seems best to place this bottom as deep as possible. This is particularly important if plants grow on the lysimeters. The bottom of a lysimeter should be so deep that only very few roots can reach it, because then the water removed by plants from the soil at the bottom will be negligible. In a rather shallow lysimeter (3 to 4 feet) many roots penetrate to the bottom layer, which is frequently better supplied with water than soil of the same depth that is in contact with strata underneath. The drier the climate, the more serious is this factor.

The movement of gravitational water through soil is not exclusively perpendicular, but frequently follows fissures, old root channels, and the like, in a slanting direction. The lysimeter walls will act as an insurmountable obstacle to such percolation. The ideal would be to construct lysimeters without side walls (Ebermayer lysimeters), but it is impossible to obtain quantitative data with them since the soil area drained is not definitely known. A practical approach to the ideal is to decrease this border effect of the side walls by building lysimeters of rather large area (one one-thousandth to one one-hundredth of an acre).

Soils containing an appreciable amount of clay have a tendency to shrink on drying, and cracks are therefore very likely to be formed between the soil block and the lysimeter walls. Precautions should be taken to avoid drainage along such cracks. Horizontal steel strips extending 1 to 2 inches into the soil and roughened side walls have been found useful in this respect. Not only will standing water alter the immediate moisture conditions of a soil, but, after some time, it will actually alter the very nature of the soil itself. It is obvious that after such a change has occurred, the data from a lysimeter lose their significance. Soils that have in nature free surface drainage should be provided with the same drainage in the lysimeter. It is obvious that the amount of run-off water should be recorded.

The writers believe that no one construction should be regarded as standard in a lysimeter and that a proper design can be made only by having an accurate knowledge of both the purpose of the experiment and of the pedologic, geologic, and climatic conditions.

EXPERIMENTAL PLAN

To avoid drawing conclusions from inadequate information, investigators should not carry out lysimeter studies without replicating each individual treatment. The number of replicates needed depends largely on the nature of the investigation and the type of lysimeters used. If the soil is left in its natural condition, no assurance is given of its uniformity; therefore, at least triplicates should be used to represent any one condition studied. The frequent practice of attempting to study too many factors with too few lysimeters should be avoided. As soil in all existing types of lysimeters may be subject to deterioration, it is necessary to continue the study of a given installation at least throughout a decade in order to determine whether such deterioration has occurred and to judge the value of the

data. It may be advisable to install a number of extra lysimeters and to remove the soil from one of them every 2 or 3 years so that its chemical, physical, and biological properties may be determined. If pronounced changes of soil properties are found, the potential future usefulness of the installation for attaining the original objective should be carefully considered.

Lysimeter installations are normally supplemented with precipitation gages and other meteorological measuring devices. For the selection of adequate equipment the investigator should get the advice of a professional meteorologist. An accurate knowledge of hydrologic factors is needed regardless of whether the main object of the investigation is the study of the rate and amount of the percolate or of its composition.

PUBLICATION OF DATA

Lysimeter investigations could be made more useful by describing the installations more completely and recording the data in a uniform manner. The soils are usually, though not always, described in detail, but information concerning soil changes during the investigation is seldom included. Any chemical, physical, and biological change should be noted, since it might give an important clue to the nature of the percolation.

All hydrologic data should be expressed as inches depth (or centimeters depth) and the chemical constituents as pounds per acre (or kilograms per hectare), so that any lysimeter investigation may be easily compared with another or with related studies. It is very confusing to read lysimeter reports in which precipitation is recorded as inches depth, percolate as cubic centimeters per lysimeter, and the chemical constituents as parts per million of the percolate.

"Lysimeter" is an accepted term for a device to study the rate, amount, and composition of natural precipitation percolating through soil and it should be used wherever reference is made to such an instrument. Such terms as "culture pot," "rim," "percolator," "percolimeter," and "drain-gauge" are ambiguous and should be avoided.

SUPPLEMENTARY STUDIES

In many instances a knowledge of the amount, rate, and composition of the percolate can well be obtained by the study of water flowing from springs or field drain tiles. This information may not be quantitatively correct, because the drainage area is not definitely known and not all the percolation water may be caught. The great value of such studies lies in the fact that the data are acquired from a percolate obtained without any interference with natural soil conditions. A combination of such studies and lysimeter research appears to be advisable.

CONCLUSION

Lysimeters have distinct limitations, as has been pointed out; but without doubt many questions concerning pedology, soil fertility, and hydrology can be answered by the correct use of lysimeters. Filled-in lysimeters may fulfill useful tasks in fertility investigations if a sound water balance is maintained in the soil; but for pedologic and hydrologic studies carefully designed monolith lysimeters seem to be indispensable.

BIBLIOGRAPHY ON LYSIMETERS

Compiled by J. M. Davidson [19]

CONTENT AND ORGANIZATION

This bibliography lists publications that refer to the design or construction of lysimeters, the method of filling and the type of soil used, the treatment and cropping of the soil, the amount of leaching and the loss of plant nutrients, the analysis of the soil and the leachate, or the discussion of some phase of lysimeter investigations. For the purpose of this bibliography a lysimeter is defined as an instrument that contains soil and receives natural rainfall or irrigation and is provided with an arrangement for collecting and measuring the percolate. Literature on natural lysimeters, such as small watersheds, having no apparent supply of moisture other than that from the atmosphere and provided with either natural drainage or tile-drainage systems, is not included. In most entries the pagination refers to that part of a publication that treats of lysimeters. No reference is included that does not deal directly with the subject of lysimeters.

The following letters are used to indicate the content of any entry:

A, Information on the size, design, construction, installation of the lysimeter.
B, Data on the amount of leaching or quantity of leachate from the lysimeter.
C, Data on the quantity of plant nutrients lost from the soil or the chemical analysis of either the leachate or the soil.
D, A comparison with other investigators' work or inclusion of data from such reports.
E, Only a general statement or discussion with no data included unless otherwise indicated.

The letter or letters directly opposite an author's name indicate those subjects discussed in the material cited in the entry. Letters within the body of the entry indicate the subject treated in particular citations.

The entries are made according to the geographical distribution of the lysimeters. The arrangement of references in chronological order under the geographical divisions indicates the historic development of lysimeter research in each. No special significance is attached to the order in which the countries are listed. Entries of the studies in the States and Territories of the United States are first; next in order are those for Great Britain, Asia, continental Europe, and Africa.

The author index (pp. 65–67) is arranged alphabetically. The numbers following each author's name provide a ready reference to the entry in the bibliography in which his work is cited.

It was the aim of the compiler of this bibliography to set up an arrangement that would permit the selection of references for any period, location, author, or specific type of subject matter without the necessity of reviewing the entire field of literature on the general subject of lysimeters. This type of bibliography reduces the time necessary for the selection of references on a particular phase of lysimeter investigation. For example, the preparation of a report on the amount of nitrogen lost by leaching from soils in lysimeters in

[19] The compiler wishes to express his gratitude for the advice and cooperation of his associates in the Soil Conservation Service, and for the excellent assistance of Robert Jones, Ann Goodwin, and others of the Works Progress Administration in the preparation of the bibliography.

Europe would necessitate the review of only those entries denoted by C and listed under European countries.

Titles of publications are given in the language in which they were published, except those in Russian. These are transliterated. If a work has appeared in more than one language the languages in which it has been published are named.

All translations by the compiler and additions to the references are in brackets. Citations to abstracts have not been made unless so stated; however, abstracts of many of the works cited appear in the Experiment Station Record and Chemical Abstracts. All the publications in this bibliography, with the exception of entry 456, which is not obtainable in this country, have been examined by the compiler in the libraries in Washington, D. C.

If omissions have been made of lysimeter investigations that are within the limitations set up for this bibliography, the librarian of the Soil Conservation Service would appreciate receiving such information. The exact title, author, publication, and date should be given. There are several lysimeter studies that have been or are being conducted in various parts of the world for which no published reports have been made.

BIBLIOGRAPHY

UNITED STATES

ARIZONA

(1) Burgess, P. S., and McGeorge, W. T. (C)
1933. A LYSIMETER STUDY OF THE NITROGEN BALANCE IN IRRIGATED ARID SOILS. Ariz. Agr. Expt. Sta. Ann. Rpt. 44: 14–15.

(2) Shreve, F. (AB)
1934. RAINFALL, RUN-OFF AND SOIL MOISTURE UNDER DESERT CONDITIONS. Assoc. Amer. Geogr. Ann. 24: 140–146, illus.

CALIFORNIA

(3) Lowdermilk, W. C. (AB)
1929. FURTHER STUDIES OF FACTORS AFFECTING SURFICIAL RUN-OFF AND EROSION. Internatl. Cong. Forestry Expt. Sta. (Stockholm 1929) Proc. pp. 606–628, illus. [Processed.]

(4) ——— (AB)
1930. INFLUENCE OF FOREST LITTER ON RUN-OFF, PERCOLATION, AND EROSION. Jour. Forestry 28: 474–491, illus.

(5) Taylor, G. H. (A)
1931. INVESTIGATIONS RELATING TO THE ABSORPTION OF PRECIPITATION AND ITS PENETRATION TO THE ZONE OF SATURATION. Natl. Res. Council, Amer. Geophys. Union Trans. 12: 206–207, illus. [Processed.]

(6) Lee, C. H. (A)
1932. ON ABSORPTION AND TRANSPIRATION. RELATION OF WATERSHED VEGETATION TO ABSORPTION. Natl. Res. Council, Amer. Geophys. Union Trans. 13: 292–293. [Processed.]

(7) American Geophysical Union, Hydrology Committee. (A)
1933. LIST OF AGENCIES IN THE UNITED STATES CARRYING ON WORK IN THE FIELD COVERED BY THE COMMITTEE ON ABSORPTION AND TRANSPIRATION, WITH BRIEF INFORMATION REGARDING THEIR WORK AND PUBLICATIONS. [FOREST SERVICE, CALIFORNIA FOREST EXPERIMENT STATION.] Natl. Res. Council, Amer. Geophys. Union Trans. 14: 357. [Processed.]

(8) Blythe, S. O. (A)
1935. WATER FOR THE YEAR 2000. A FIFTY YEAR EXPERIMENT AT SAN DIMAS FOR INDUSTRY AND AGRICULTURE. Calif. Mag. Pacific Business 26 (11): 20–23, 50–52, illus.

(9) Kraebel, C. J. (E)
1935. run-off and erosion experiments in mountain areas. plot and lysimeter studies. Amer. Soc. Civ. Engin. Water Conserv. Prog. Rpt., Los Angeles, 1935: 92–94. [Mimeographed.]
(10) California Forest and Range Experiment Station. (A)
1936. lysimeters. U. S. Forest Serv., San Dimas Expt. Forest. (Ed. 1935, rev.) pp. 7, 19, illus.

CONNECTICUT

(11) Anderson, P. J., Swanback, T. R., and Street, O. E. (E)
1930. report of the tobacco substation for 1929. Conn. (State) Agr. Expt. Sta. Bul. 311: 205–206, illus.
(12) Connecticut Agricultural Experiment Station. (A)
1930. soils. leaching experiments. Conn. (State) Agr. Expt. Sta. (Ann. Rpt. 1929) Bul. 318: 762–763, illus.
(13) ——— (BC)
1931. nitrogen loss revealed by lysimeter. Conn. (State) Agr. Expt. Sta. (Ann. Rpt. 1930) Bul. 322: 135–136.
(14) Morgan, M. F., Street, O. E., and Jacobson, H. G. M. (ABC)
1931. fertilizer losses through leaching as measured by lysimeter experiments. Conn. (State) Agr. Expt. Sta. (Tobacco Sta. Rpt. 1930) Bul. 326: 432–441.
(15) Connecticut Agricultural Experiment Station. (AC)
1932. Lysimeters show top-dressing necessary. Conn. (State) Agr. Expt. Sta. (Ann. Rpt. 1931) Bul. 337: 477–479.
(16) ——— (C)
1933. soils. Conn. (State) Agr. Expt. Sta. (Ann. Rpt. 1932) Bul. 347: 280.
(17) ——— (C)
1934. soils. lysimeter experiments. Conn. (State) Agr. Expt. Sta. (Ann. Rpt. 1933) Bul. 357: 131–133.
(18) ——— (BC)
1935. lysimeter data from forest soils. Conn. (State) Agr. Expt. Sta. (Ann. Rpt. 1934) Bul. 366: 90–91, 97.
(19) Lunt, H. A. (ABC)
1935. forest lysimeter studies under pine. Amer. Soil Survey Assoc. Bul. 16: 86–92, illus. [Mimeographed.]
(20) Connecticut Agricultural Experiment Station. (BC)
1936. drainage losses under dolomitic lime treatment. forest lysimeters. Conn. (State) Agr. Expt. Sta. (Ann. Rpt. 1935) Bul. 381: 190–191.
(21) Morgan, M. F. (ACD)
1936. soil changes resulting from nitrogeneous fertilization. a lysimeter study. Conn. (State) Agr. Expt. Sta. Bul. 384: 371–449, illus.
(22) Connecticut Agricultural Experiment Station. (E)
1937. nitrogeneous fertilizer and soil acidity. Conn. (State) Agr. Expt. Sta. (Ann. Rpt. 1936) Bul. 393: 195–196.
(23) Lunt, H. A. (ABCD)
1937. forest lysimeter studies under red pine. Conn. (State) Agr. Expt. Sta. Bul. 394: 221–268, illus.
(24) Morgan, M. F., and Bailey, E. M. (E)
1937. the evaluation of the influence of nitrogeneous fertilizers on the acid-base status of soils by lysimeter studies. Amer. Fert. 87 (5): 7–8.

FLORIDA

(25) Blair, A. W. (A)
1911. soil tank investigations Fla. Agr. Expt. Sta. Ann. Rpt. 1910: 32–34, illus.
(26) ——— and Collison, S. E. (A)
1911. tanks for soil investigations at the florida agricultural experiment station. Jour. Indus. and Engin. Chem. 3: 32–33. illus.

(27) BLAIR, A. W. (AB)
1911. LOSS OF FERTILIZER CONSTITUENTS IN DRAINAGE WATER. Fla. State Hort. Soc. Proc. 24: 105–111, illus.

(28) ——— (BC)
1912. SOIL TANK INVESTIGATIONS. Fla. Agr. Expt. Sta. Ann. Rpt. 1911: 32–39.

(29) COLLISON, S. E. (BC)
1913–16. SOIL TANK INVESTIGATIONS. Fla. Agr. Expt. Sta. Ann. Rpts. 1912: 43–47; 1915: 102–105.

(30) ——— (BC)
1914–15. SOIL TANK EXPERIMENTS. Fla. Agr. Expt. Sta. Ann. Rpts. 1913: 98–102; 1914: 77–80.

(31) ——— and WALKER, S. S. (BC)
1916. LOSS OF FERTILIZERS BY LEACHING. Fla. Agr. Expt. Sta. Bul. 132, 20 pp., illus.

(32) ——— (C)
1917–19. SOIL TANK INVESTIGATIONS. Fla. Agr. Expt. Sta. Ann. Rpts. 1916: 116R–118R; 1917: 101R–102R; 1919: 43R–45R.

(33) ——— (AC)
1919. CITRUS FERTILIZER EXPERIMENTS. COMPOSITION OF SOIL. LEACHING OF FERTILIZER. Fla. Agr. Expt. Sta. Bul. 154: 7–15, illus.

(34) RUPRECHT, R. W. (E)
1921. REPORT OF PHYSIOLOGICAL CHEMIST. Fla. Agr. Expt. Sta. Ann. Rpt. 1921: 22R–24R.

(35) ——— (E)
1922. CITRUS PROBLEMS. Fla. Agr. Expt. Sta. Ann. Rpt. 1922: 42R–43R.

(36) ——— (BC)
1923–25. DIEBACK OF CITRUS. Fla. Agr. Expt. Sta. Ann. Rpts. 1923: 46R–48R; 1924: 47R–49R; 1925: 28R–30R.

(37) ——— (E)
1926–28. DIEBACK OF CITRUS. SOIL STUDIES. GREEN MANURE PROJECT. Fla. Agr. Expt. Sta. Ann. Rpts. 1926, Dieback of Citrus, pp. 29R–30R; 1927, Soil Studies, pp. 33R–34R; 1928, Green Manure Project, p. 34R.

(38) ——— (BC)
1929. DETERMINATION OF THE EFFECT OF GREEN MANURE ON THE COMPOSITION OF THE SOIL. LYSIMETER STUDIES. Fla. Agr. Expt. Sta. Ann. Rpt. 1929: 47–48.

(39) STOKES, W. E. (E)
1930. LYSIMETER STUDIES ON PASTURE GRASSES. GREEN MANURE STUDIES. Fla. Agr. Expt. Sta. Ann. Rpt. 1930: 36, 47.

(40) RUPRECHT, R. W. (E)
1930. DETERMINATION OF THE EFFECT OF GREEN MANURES ON COMPOSITION OF SOIL. Fla. Agr. Expt. Sta. Ann. Rpt. 1930: 60–61.

(41) BARNETTE, R. M. (CE)
1931–34. DETERMINATION OF THE EFFECT OF GREEN MANURES ON THE COMPOSITION OF SOIL. LYSIMETER EXPERIMENTS. Fla. Agr. Expt. Sta. Ann. Rpts. 1931: 63–64 [C]; 1932: 65–66 [E]; and 1934: 49 [E].

(42) ——— (AC)
1933. DETERMINATION OF THE EFFECT OF GREEN MANURES ON THE COMPOSITION OF SOIL. A STUDY OF THE DECOMPOSITION OF FOREST, RANGE, AND PASTURE GROWTHS TO FORM SOIL ORGANIC MATTER. Fla. Agr. Expt. Sta. Ann. Rpt. 1933: 71–74.

(43) ——— JONES, H. W., and HESTER, J. B. (ABC)
1938. LYSIMETER STUDIES WITH THE DECOMPOSITION OF SUMMER COVER CROPS. Fla. Agr. Expt. Sta. Bul. 327, 44 pp., illus.

(44) BENSON, N., and BARNETTE, R. M. (AC)
1939. LEACHING STUDIES WITH VARIOUS SOURCES OF NITROGEN. Amer. Soc. Agron. Jour. 31: 44–54.

HAWAII

(45) Eckart, C. F. (C)
1901. report on fertilization. Hawaii. Sugar Planters' Assoc. Fert. Comn. Rpt. 1901: 14–24.

(46) ——— (ABC)
1902. plan of experiment. fertilization of lysimeter soils. work of the experiment station and laboratories. Hawaii. Sugar Planters' Assoc. Work Rpt. 1902: 26–100, illus.

(47) Maxwell, W. (D)
1905. statement of the work of the experiment station and laboratories. a lysimeter. Hawaii. Sugar Planters' Assoc. Expt. Sta. (Div. Agr. and Chem. Ann. Rpt. 1897) Bul. 3: 6.

(48) ——— (ABC)
1905. report on fertilization, notes on nitrogen, lime, and irrigation. Hawaii. Sugar Planters' Assoc. Expt. Sta. (Div. Agr. and Chem. Ann. Rpt. 1898) Bul. 4: 12–14, 20–24.

(49) ——— (D)
1905. work of the experiment station and laboratories. some results of over-irrigation. Hawaii. Sugar Planters' Assoc. Expt. Sta. (Div. Agr. and Chem. Ann. Rpt. 1899) Bul. 5: 21.

(50) Eckart, C. F. (ABC)
1905. a consideration of the action of saline irrigation water; the action of salt on the soil. plan of experiments. Hawaii. Sugar Planters' Assoc. Expt. Sta. (Div. Agr. and Chem. Ann. Rpt. 1902) Bul. 8: 23–74. illus.

(51) ——— (BC)
1906. lysimeter experiments. Hawaii. Sugar Planters' Assoc. Expt. Sta. (Div. Agr. and Chem.) Bul. 19, 31 pp.

(52) Thompson, Alice R. (C)
1909. report of the assistant chemist. water analysis. Hawaii. Agr. Expt. Sta. Ann. Rpt. 1908: 62.

(53) Eckart, C. F. (E)
1910. report of committee in charge of experiment station. research work in hand. Hawaii. Planters' Rec. 3: 303–304.

(54) ——— (E)
1911. report of the experiment station committee. chemistry and sugar technology research. lysimeter experiments. Hawaii. Planters' Rec. 5: 335.

(55) Peck, S. S. (AC)
1911. lysimeter experiments. Hawaii. Sugar Planters' Assoc. Expt. Sta. (Agr. and Chem. Ser.) Bul. 37, 38 pp., illus.

(56) ——— (AC)
1912. the influence of molasses on nitrification of cane soils. Hawaii. Sugar Planters' Assoc. Expt. Sta. (Agr. and Chem. Ser.) Bul. 39, 27 pp., illus.

(57) ——— (AC)
1914. the influence of fineness upon the availability of bone meal. lysimeter tests. Hawaii. Planters' Rec. 10: 419-425.

(58) ——— (C)
1915. lysimeter experiments with press cake. Hawaii. Planters' Rec. 12: 207-211.

ILLINOIS

(59) Stauffer, R. S., and Smith, R. S. (ABD)
1937. variation in soils with respect to the dispostion of natural precipitation. Amer. Soc. Agron. Jour. 29: 917–923, illus.

INDIANA

(60) Indiana Agricultural Experiment Station. (AB)
1890. investigations undertaken. condition of uncompleted work. Ind. Agr. Expt. Sta. Ann. Rpt. (1889) 2: 21–22, 31–33.

(61) Huston, H. A., and Goss, A. (C)
1890. the absorptive power of soils. adsorption exercised by soil on purdue university farm. Ind. Agr. Expt. Sta. Bul. 33: 50–54.

(62) WILEY, H. W. (AE)
1894. PRINCIPLES AND PRACTICE OF AGRICULTURAL ANALYSIS. I (SOILS). 606 pp., illus. Easton, Pa. *See* Lysimetry, ed. 1, p. 158; ed. 2, p. 173; ed. 3, p. 186 [E]; and Measurement of Percolation Through the Soil in Situ, ed. 1, pp. 164–167; ed. 2, pp. 176–178; ed. 3, pp. 191–194 [A].

IOWA

(63) MUSGRAVE, G. W. (E)
1934. A QUANTITATIVE STUDY OF CERTAIN FACTORS AFFECTING SOIL- AND WATER-LOSSES AS THE LOGICAL BASIS FOR DEVELOPING PRACTICAL METHODS OF EROSION CONTROL. Natl. Res. Council, Amer. Geophys. Union Trans. 15 (pt. 2): 515–521. [Processed.]

(64) ——— (E)
1935. THE INFILTRATION CAPACITY OF SOILS IN RELATION TO THE CONTROL OF SURFACE RUN-OFF AND EROSION. Amer. Soc. Agron. Jour. 27: 336–345, illus.

(65) ——— (ABD)
1935. A DEVICE FOR MEASURING PRECIPITATION WATERS LOST FROM THE SOIL AS SURFACE RUN-OFF, PERCOLATION, EVAPORATION, AND TRANSPIRATION. Soil Sci. 40: 391–401, illus.

(66) AYRES, Q. C. (A)
1936. SOIL EROSION AND ITS CONTROL. 365 pp., illus. New York. *See* Absorption Range of Soils, pp. 40–42 [A].

(67) MUSGRAVE, G. W., and NORTON, R. A. (B)
1937. SOIL AND WATER CONSERVATION INVESTIGATIONS AT THE SOIL CONSERVATION EXPERIMENT STATION, MISSOURI VALLEY LOESS REGION, CLARINDA, IOWA, PROGRESS REPORT, 1931–35. U S. Dept. Agr. Tech. Bul. 558, 181 pp., illus. *See* Purpose and Plan of Experiments, pp. 14–20, Lysimeter Measurements of Percolate, Run-off, and Erosion, pp. 37–41; and Individual Run-off and Monthly Percolation Records of Lysimeters, pp. 144–158.

(68) ——— and NEAL, O. R. (B)
1937. RAINFALL AND RELATIVE LOSSES IN VARIOUS FORMS. Natl. Res. Council, Amer. Geophys. Union Trans. 18 (pt. 2): 349–355, illus. [Processed.]

(69) NEAL, O. R., RICHARDS, L. A., and RUSSELL, M. B. (AB)
1938. OBSERVATIONS ON MOISTURE CONDITIONS IN LYSIMETERS. Soil Sci. Soc. Amer. Proc. (1937) 2: 35–44, illus. [Processed.]

KENTUCKY

(70) KARRAKER, P. E. (A)
1934. AN INEXPENSIVE LYSIMETER. Amer. Soc. Agron. Jour. 26: 75, illus.

(71) ——— and BORTNER, C. E. (AC)
1938. NITROGEN LEACHING IN SOIL ON THE EXPERIMENT STATION FARM AT LEXINGTON. Soil Sci. Soc. Amer. Proc., Chicago, 1937, 2:393–398. [Processed.]

(72) ——— (BC)
1939. NITROGEN REMOVED BY DRAINAGE AND CROPPING FROM LYSIMETERS AS AFFECTED BY KIND OF VEGETATIVE COVER. Com. Fert. 58 (3): 18–19, illus.

MASSACHUSETTS

(73) STURTEVANT, E. L. (A)
1876. WAUSHAKUM FARM EXPERIMENTS. 3. LYSIMETER. Sci. Farmer 1: 90.

(74) ——— (BC)
1876. WAUSHAKUM FARM EXPERIMENTS. LYSIMETERS. Sci. Farmer 1: 125–126, 176.

(75) ——— (BD)
1877. WAUSHAKUM FARM NOTES. LYSIMETERS. Sci. Farmer 2: 16, 30, 100.

(76) ——— (ABD)
1878. WAUSHAKUM FARM CONTRIBUTIONS. LYSIMETERS OR DIFFERENTIAL RAIN GAUGES. A NEW LYSIMETER. ANOTHER LYSIMETER. Sci. Farmer 3, Waushakum Farm Contributions, pp. 17–18 [AB]; Lysimeters or Differential Rain Gauges, pp. 49–50, illus. [D]; A New Lysimeter, p. 82 [A]; and Another Lysimeter, p. 97 [A].

(77) STURTEVANT, E. L. (A)
1879. ABSORPTIVE POWER OF SOILS. (LYSIMETER). Mass. Bd. Agr. Sec. Ann. Rpt. (1878) 26: 71–72.
(78) ——— (AB)
1879. MASSACHUSETTS AGRICULTURAL COLLEGE EXPERIMENT STATION LYSIMETER. LYSIMETER RESULTS. Sci. Farmer 4: 59, 101.
(79) STOCKBRIDGE, L. (AB)
1879. METHODS OF CONSTRUCTING THE LYSIMETER. Mass. Agr. Col. Ann. Rpt. 16: 45–51, illus.
(80) GOESSMANN, C. A. (C)
1879. REPORT ON THE PERCOLATED WATERS OF THE LYSIMETER. Mass. Agr. Col. Ann. Rpt. 16: 51–54.
(81) STORER, F. H. (BD)
1888–97. AGRICULTURE IN SOME OF ITS RELATIONS WITH CHEMISTRY. Ed. [1], v. 1, New York, 1888; ed. 7, 620 pp., 1897. *See* Drain Gauges, ed. [1], pp. 62–68 [BD]; and American Experience, ed. 7, pp. 88–90 [BD].
(82) ANONYMOUS. (ABD)
1915. LYSIMETER OR DRAIN GAUGE. *In* New International Encyclopaedia, ed. 2, v. 14, pp. 523–524, illus. New York.
(83) THAYER, C. H. (E)
1936. STOCKBRIDGE, LEVI. *In* Dictionary American Biographies, v. 18, pp. 38–39. New York.
(84) EDWARDS, E. E. (E)
1936. STURTEVANT, EDWARD LEWIS. *In* Dictionary American Biographies, v. 18, pp. 185–186. New York.

MISSOURI

(85) SANBORN, J. W. (AB)
1889. A NEW LYSIMETER. Soc. Prom. Agr. Sci. Proc. 10: 54–59, illus.
(86) MILLER, M. F., and DULEY, F. L. (A)
1917–19. STUDIES OF WATER ABSORPTION, PERCOLATION, EVAPORATION, CAPILLARY WATER MOVEMENT AND EROSION UNDER FIELD CONDITIONS. Mo. Agr. Expt. Sta. (Ann. Rpts. 1917, 1918) Bul. 151: 58–59; Bul. 163: 65–66.
(87) DULEY, F. L. (A)
1925. A MOVABLE LYSIMETER FOR SOIL STUDIES. Soil Sci. 20: 465–471, illus.

NEW JERSEY

(88) JOFFE, J. S. (A)
1929. A NEW TYPE OF LYSIMETER AT THE NEW JERSEY AGRICULTURAL EXPERIMENT STATION. Science 70: 147–148, illus.
(89) ——— (ABD)
1932. LYSIMETER STUDIES I. MOISTURE PERCOLATION THROUGH THE SOIL PROFILE. Soil Sci. 34: 123–143, illus.
(90) ——— (BCD)
1933. LYSIMETER STUDIES II. THE MOVEMENT AND TRANSLOCATION OF SOIL CONSTITUENTS IN THE SOIL PROFILE. Soil Sci. 35: 239–257, illus.
(91) ——— (BCD)
1933. LYSIMETER STUDIES III. THE MOVEMENT AND TRANSLOCATION OF NITROGEN AND ORGANIC CONSTITUENTS IN THE PROFILE OF A PODZOLIC SOIL. Soil Sci. 35: 401–411.
(92) ——— (E)
1936. PEDOLOGY. 575 pp., illus. New Brunswick, N. J. *See* Prairie Soils, Subtypes and Transition Types of Soils in Podsol Zone, and The Yellow Soils, pp. 229, 231–232, 301, 413.

NEW YORK

(93) NEW YORK STATE AGRICULTURAL EXPERIMENT STATION. (AB)
1883. LYSIMETERS OR DRAIN GAUGES. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1882) 1: 14–16.
(94) ——— (B)
1884. LYSIMETER. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1883) 2: 31.
(95) BABCOCK, S. M. (BCD)
1885. LYSIMETERS. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1884) 3: 329, 347–350.

(96) Babcock, S. M. (BC)
1886–88. lysimeters. N. Y. State Agr. Expt. Sta. Ann. Rpts. (1885) 4: 293, 321–325; (1886) 5: 326–328; (1887) 6: 389, 393–398.

(97) Goff, E. S. (A)
1888. eine studie über die bewegungen des bodenwasser. ein neues lysimeter. Wollny's Agr.-Phys. 11: 71–72, illus.

(98) ——— (A)
1888. a new lysimeter. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1887) 6: 113–114, illus.

(99) ——— (A)
1889. details of the construction of new lysimeters. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1888) 7: 187–191, illus.

(100) Ladd, E. F. (BC)
1889. amount and analysis of drainage from each lysimeter for 1888. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1888) 7: 313–314.

(101) Emery, F. E., and Goff, E. S. (B)
1891. report on lysimeters for 1889. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1890) 9: 352, 390–401.

(102) Warington, R. (BD)
1900. lectures on the physical properties of soil. 231 pp., illus. Oxford, England. *See* Influence of a Crop on Evaporation, pp. 122–124.

(103) Horton, R. E. (AB)
1907–11. græfenburg hydrophysical station near utica, n. y. N. Y. State Engin. and Surveyor Ann. Rpts. Sup. 1906: 215–245, illus.; 1907: 518–532; 1910: 552–591.

(104) Lyon, T. L. (A)
1909. tanks for soil investigation at cornell university. Science (n. s.) 29: 621–623, illus.

(105) New York Agricultural Experiment Station. (E)
1915. soil investigations. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1914) 33: 19.

(106) Lyon, T. L., Fippin, E. O., and Buckman, H. O. (ACD)
1915. soils, their properties and management. 764 pp., illus. New York. *See* The Lysimeter Method of Studying Gravitational Water, pp. 240–241, illus. [AD]; Losses in Nitrogen and Calcium Composition of Surface Water, pp. 372–374 [C].

(107) New York State Agricultural Experiment Station. (E)
1916. investigations guide to buildings and grounds. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1915) 34: 25, 973.

(108) Barker, J. F. (A)
1917–18. facilities for lysimeter and outdoor pot culture work at the station. N. Y. State Agr. Expt. Sta. Tech. Bul. 61, 10 pp., illus. Reprinted in Ann. Rpt. (1917) 36: 47–54, illus.

(109) Lyon, T. L., and Bizzell, J. A. (ABCD)
1918. lysimeter experiments. records for tanks 1 to 13 during the years 1910–1914, inclusive. N. Y. (Cornell) Agr. Expt. Sta. Mem. 12, 115 pp., illus.

(110) New York State Agricultural Experiment Station. (E)
1919. results of stations work in 1918. lysimeter work. laboratory work. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1918) 37: 11–12.

(111) Hedrick, U. P. (A)
1919. sturtevant's notes on edible plants. N. Y. State Dept. Agr. Ann. Rpt. 27 (pt. 2): 4.

(112) Lyon, T. L., and Bizzell, J. A. (BC)
1921. lysimeter experiments ii. records for tanks 13 to 16 during the years 1913 to 1917 inclusive. N. Y. (Cornell) Agr. Expt. Sta. Mem. 41: 45–93.

(113) New York State Agricultural Experiment Station. (E)
1922–31. lysimeter investigations. N. Y. State Agr. Expt. Sta. Ann. Rpts. (1921) 40: 17; (1923) 42: 25; (1924) 43: 20; (1926) 45: 16–17; (1927) 46: 22; (1929) 48: 24; (1930) 49: 25, 80.

(114) Bizzell, J. A. (C)
1922. disappearance of nitrates from soil under timothy. nitrates removed by leaching. Amer. Soc. Agron. Jour. 14: 324–325.

(115) Lyon, T. Lyttleton, and Buckman, Harry O. (ABCD)
1923–37. THE NATURE AND PROPERTIES OF SOILS; A COLLEGE TEXT OF EDAPHOLOGY. Ed. 1, 588 pp., illus. New York; rev. ed. [ed. 2] 428 pp., illus., 1929; ed. 3, 392 pp., illus., 1937. *See* Importance of the Study of the Flow and Composition of Drainage Water, ed. 1, pp. 178–182 [AD]; Percolation Losses and their Control, ed. 1, pp. 206–210 [BCD]; Quantities of Nutrients Removed by Drainage and Cropping, Possible Exhaustion of the Soil, ed. 1, pp. 307–310 [C]; Percolation Losses and their Control, rev. ed., pp. 214–218 [BD]; Methods of Studying Drainage Water, and Amounts of Nutrients Removed from Soil by Drainage, rev. ed., pp. 249–257 [ACD]; Changes of Lime in Soil-Losses, rev. ed., pp. 289–290 [C]; The Gravitational Water, ed. 3, pp. 119–120 [C]; Percolation Methods of Study, and Percolation Losses of Nutrients, ed. 3, pp. 146–153, illus. [ABCD]; and Changes of Lime in the Soil and the Subsequent Losses, ed. 3, pp. 266–268 [C].

(116) New York State Agricultural Experiment Station. (C)
1923–26. LYSIMETER INVESTIGATIONS. N. Y. State Agr. Expt. Sta. Ann. Rpts. (1922) 41: 23; (1925) 44: 19.

(117) Wilson, B. D. (BCD)
1923. THE EFFECT OF PLANTS ON THE CONCENTRATION OF DRAINAGE WATER FROM THE CORNELL LYSIMETERS. Soil Sci. 16: 427–432.

(118) Bizzell, J. A. (CD)
1926. REMOVAL OF PLANT NUTRIENTS IN DRAINAGE WATERS. Amer. Soc. Agron. Jour. 18: 130–136.

(119) Barrows, H. K. (ABD)
1927. WATER POWER ENGINEERING. 734 pp., illus. New York and London. *See* Measurements of Soil Evaporation, pp. 91–95.

(120) Bizzell, J. A., and Lyon, T. L. (ABC)
1928. COMPOSITION OF DRAINAGE WATERS FROM LYSIMETERS AT CORNELL UNIVERSITY. 1st Internatl. Cong. Soil Sci., Washington, D. C., 1927, Proc. and Papers, Comn. 2, v. 2, pp. 342–349.

(121) New York State Agricultural Experiment Station. (CE)
1929. LYSIMETER INVESTIGATIONS. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1928) 47: 18–19.

(122) Collison, R. C., and Mensching, J. E. (ABCD)
1930. LYSIMETER INVESTIGATIONS I. NITROGEN AND WATER RELATIONS OF CROPS IN LEGUME AND NON-LEGUME ROTATIONS. N. Y. State Agr. Expt. Sta. Tech. Bul. 166, 90 pp., illus.

(123) Lyon, T. L., Bizzell, J. A., Wilson, B. D., and Leland, E. W. (BC)
1930. LYSIMETER EXPERIMENTS III. RECORD FOR TANKS 3 TO 12 DURING THE YEARS 1910 TO 1924 INCLUSIVE. N. Y. (Cornell) Agr. Expt. Sta. Mem. 134, 72 pp.

(124) New York State Agricultural Experiment Station. (C)
1932. NITROGEN AND WATER RELATIONS IN LEGUME AND NON-LEGUME CROP ROTATIONS. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1931) 50: 74.

(125) Collison, R. C., and Mensching, J. E. (CD)
1932. LYSIMETER INVESTIGATIONS II. COMPOSITION OF RAINWATER AT GENEVA, N. Y., FOR A 10 YEAR PERIOD. N. Y. State Agr. Expt. Sta. Tech. Bul. 193, 19 pp.

(126) New York State Agricultural Experiment Station. (C)
1933. NEW METHODS FOR STUDYING ORCHARD SOILS: MINERAL RELATIONS IN LEGUME AND NON-LEGUME ROTATIONS. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1932) 51: 86–87, 89.

(127) Lyon, T. L., and Bizzell, J. A. (AC)
1933. LYSIMETER EXPERIMENTS WITH SULPHATE OF AMMONIA AND NITRATE OF SODA. Jour. Agr. Res. 47: 53–63.

(128) Collison, R. C., Beattie, H. G., and Harlan, J. D. (BC)
1933. LYSIMETER INVESTIGATIONS III. MINERAL AND WATER RELATIONS AND FINAL NITROGEN BALANCE IN LEGUME AND NON-LEGUME CROP ROTATIONS FOR A PERIOD OF 16 YEARS. N. Y. State Agr. Expt. Sta. Tech. Bul. 212, 81 pp. illus.

(129) New York State Agricultural Experiment Station. (E)
1934. TREE NUTRITION. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1933) 52: 82.

(130) NEW YORK STATE AGRICULTURAL EXPERIMENT STATION. (E)
1935. LYSIMETERS SHOW MANY FUNDAMENTAL SOIL RELATIONS. TREE RESPONSES STUDIED BY LYSIMETER METHODS. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1934) 53: 60–61.

(131) COLLISON, R. C. (ABC)
1935. LYSIMETER INVESTIGATIONS IV. WATER MOVEMENT, SOIL TEMPERATURES, AND ROOT ACTIVITY UNDER APPLE TREES. N. Y. State Agr. Expt. Sta. Tech. Bul. 237, 31 pp.

(132) NEW YORK STATE AGRICULTURAL EXPERIMENT STATION. (C)
1936. LYSIMETER INVESTIGATIONS WITH TWO TYPES OF EQUIPMENT. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1935) 54: 16, 78.

(133) LYON, T. L., and BIZZELL, J. A. (BC)
1936. LYSIMETER EXPERIMENTS IV. RECORDS FOR TANKS 17 TO 20 DURING THE YEARS 1922 TO 1933, AND FOR TANKS 13 TO 16 DURING THE YEARS 1913 TO 1928. N. Y. (Cornell) Agr. Expt. Sta. Mem. 194, 59 pp.

(134) NEW YORK STATE AGRICULTURAL EXPERIMENT STATION. (E)
1937. NUTRIENT AND SOIL MOISTURE STUDIES WITH LYSIMETERS. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1936) 55: 67–68.

(135) ——— (E)
1938. RENEWED INTEREST IN THE USE OF POTASH IN ORCHARDS: WATER, NUTRIENT, AND CULTURAL RELATIONS OF APPLE TREES AND ROOT STUDIES. N. Y. State Agr. Expt. Sta. Ann. Rpt. (1937) 56: 67–68.

(136) UTZ, E. J. (CD)
1938. THE NATURE AND EXTENT OF SOIL LOSSES. SOIL LOSSES THROUGH LEACHING. U. S. Dept. Agr. Yearbook (Soils and Men) 1938: 87.

NORTH CAROLINA

(137) BARTEL, F. O., and SLATER, C. S. (B)
1938. PROGRESS REPORT OF THE CENTRAL PIEDMONT SOIL AND WATER CONSERVATION EXPERIMENT STATION, STATESVILLE, N. C. 1930–1935: EQUIPMENT AND METHODS OF INVESTIGATIONS; LYSIMETERS; LYSIMETER STUDIES. U. S. Soil Conserv. Serv. ESR–6: 30, 46. [Mimeographed.]

OHIO

(138) GARSTKA, W. U. (A)
1937. IN-PLACE LYSIMETERS NEAR COSHOCTON, OHIO. U. S. Soil Conserv. Serv., Soil Conserv. 3: 91–92, illus.

(139) ——— (A)
1938. DESIGN OF THE AUTOMATIC RECORDING IN-PLACE LYSIMETERS NEAR COSHOCTON, OHIO. Soil Sci. Soc. Amer. Proc. (1937) 2: 555–559, illus. [Processed.]

(140) RIESBOL, H. S., and SHERMAN, G. L. (A)
1938. WATERCYCLE LYSIMETERS FOR WATERSHED STUDIES. Agr. Engin. 19: 123–128, illus.

OREGON

(141) DEAN, H. K. (AB)
1917. THE WORK OF THE UMATILLA RECLAMATION PROJECT EXPERIMENT FARM IN 1915 AND 1916: LYSIMETER INVESTIGATIONS. U. S. Bur. Plant Indus., West. Irrig. Assoc. Cir. 17: 14–16.

(142) ——— (AB)
1919. THE WORK OF THE UMATILIA RECLAMATION PROJECT EXPERIMENT FARM IN 1917: LYSIMETER INVESTIGATIONS. U. S. Bur. Plant Indus., West. Irrig. Assoc. Cir. 26: 18–20.

(143) ——— (ABC)
1920. THE WORK OF THE UMATILLA RECLAMATION PROJECT EXPERIMENT FARM IN 1918 AND 1919: LYSIMETER INVESTIGATIONS. U. S. Dept. Agr. Dept. Cir. 110: 19–21, illus.

(144) ——— (BC)
1925. WORK OF THE UMATILLA RECLAMATION PROJECT EXPERIMENT FARM IN 1920, 1921, AND 1922: LYSIMETER INVESTIGATIONS. U. S. Dept. Agr. Dept. Cir. 342: 21–22.

(145) Dean, H. K. (AB)
1927. work of the umatilla field station in 1923, 1924, and 1925: lysimeter investigations. U. S. Dept. Agr. Dept. Cir. 422: 14–15.
(146) Higby, W. M. (CD)
1927. lysimeter studies. Soil Sci. 24: 51–56.

PUERTO RICO

(147) Hendrickson, H. C. (C)
1929. citrus investigations: pot and lysimeter experiments. Puerto Rico Agr. Expt. Sta. Ann. Rpt. 1928: 28–29.

SOUTH CAROLINA

(148) Adams, J. E. (A)
1933. green manuring and fertilizer studies in lysimeters. S. C. Agr. Expt. Sta. Ann. Rpt. 46: 154–156, illus.
(149) ——— (AC)
1934. soil fertility studies: green manuring and fertilizer studies in lysimeters. S. C. Agr. Expt. Sta. Ann. Rpt. 47: 128–130.
(150) McKaig, N., Jr. (BC)
1935. soil fertility studies: lysimeter investigations. S. C. Agr. Expt. Sta. Ann. Rpt. 48: 130–131.
(151) ——— and Roller, E. M. (BC)
1936–37. soil fertility studies: lysimeter investigations. S. C. Agr. Expt. Sta. Ann. Rpts. 49: 115–119; 50: 130–135, illus.
(151a) ——— and Roller, E. M. (AC)
1939. the effects of organic matter added to lysimeters containing norfolk coarse sand. Soil Sci. Soc. Amer. Proc. (1938) 3: 195–204, illus. [Processed.]

TENNESSEE

(152) Mooers, C. A., and MacIntire, W. H. (A)
1915. two equipments for investigation of soil leachings. a pit equipment. a hillside equipment. Tenn. Agr. Expt. Sta. Bul. 111, 8 pp., illus.
(153) ——— (AC)
1917. a comparative study of the nitrogen economy of certain tennessee soils. Tenn. Agr. Expt. Sta. Bul. 118: 127–187, illus.
(154) MacIntire, W. H., Willis, L. G., and Holding, W. A. (C)
1917. the divergent effects of lime and magnesia upon the conservation of soil sulfur. Soil Sci. 4: 231–237, illus.
(155) Mooers, C. A. (AE)
1919. adams fund projects. Tenn. Agr. Expt. Sta. Ann. Rpt. (1918) 31: 10–12.
(156) MacIntire, W. H. (ACD)
1919. the carbonation of burnt lime in soils. Soil Sci. 7: 325–453, illus.
(157) ——— (CD)
1919. the liberation of native soil potassium induced by different calcic and magnesic materials, as measured by lysimeter leachings. Soil Sci. 8: 337–395, illus.
(158) ——— and Mooers, C. A. (A)
1921. a pitless lysimeter equipment. Soil Sci. 11: 207–213, illus.
(159) ——— and Young, J. B. (C)
1923. sulfur, calcium, magnesium, and potassium content and reaction of rainfall at different points in tennessee. Soil Sci. 15: 205–227.
(160) ——— Shaw, W. M., and Young, J. B. (BCD)
1923. the influence of calcic and magnesic additions upon the outgo of sulfates from a loam soil as measured by lysimeter leachings over an 8-year period. Soil Sci. 16: 1–40; illus.

(161) MacIntire, W. H., Shaw, W. M., and Young, J. B. (BC)
1923. INFLUENCE OF CALCIC AND MAGNESIC TREATMENTS UPON THE SULFATE LEACHINGS FROM SOIL ALONE AND WITH ADDITIONS OF FERROUS SULFATE, PYRITE AND SULFUR. Soil Sci. 16: 159–182, illus.

(162) ——— Shaw, W. M., and Young, J. B. (ACD)
1923. A FIVE YEAR LYSIMETER STUDY OF THE SUPPOSED LIBERATION OF SOIL POTASSIUM BY CALCIC AND MAGNESIC ADDITIONS. Soil Sci. 16: 217–223, illus.

(163) ——— Shaw, W. M., and Young, J. B. (C)
1923. THE VARIANT ROLES OF SOIL AND SUBSOIL IN CALCIUM-MAGNESIUM INTERCHANGE. Soil Sci. 16: 321–341.

(164) ——— Shaw, W. M., and Young, J. B. (C)
1923. RECIPROCAL REPRESSION EXERTED BY CALCIC AND MAGNESIC ADDITIONS UPON THE SOLUBILITY OF NATIVE MATERIALS IN SURFACE SOIL. Soil Sci. 16: 449–464.

(165) ——— and Shaw, W. M. (CD)
1924. THE EFFECT OF SOIL SUSPENSIONS UPON THE SOLUBILITY OF THE SULFATE RADICAL IN THE SYSTEM, $Ca(OH)_2$–$CaSO_4$–H_2O. Soil Sci. 17: 65–89.

(166) ——— and Young, J. B. (CD)
1925. NITRATE LEACHINGS AS INFLUENCED BY CALCIC AND MAGNESIC ADDITIONS, WITH AND WITHOUT SULFUR CARRIERS. Soil Sci. 19: 309–323, illus.

(167) ——— and Shaw, W. M. (ACD)
1925. THE DISINTEGRATION OF LIMESTONE AND DOLOMITE SEPARATES AS INFLUENCED BY ZONE OF INCORPORATION. Soil Sci. 20: 403–417, illus.

(168) ——— Shaw, W. M., Young, J. B., and Robinson, B. (C)
1926. THE EFFECTS OF 12-YEAR RESIDUES OF LIME AND MAGNESIA UPON THE OUTGO OF SUBSEQUENT ADDITIONS OF POTASH. Amer. Soc. Agron. Jour. 18: 202–215, illus.

(169) ——— (C)
1926. INFLUENCE OF FORM, SOIL-ZONE, AND FINENESS OF LIME AND MAGNESIA INCORPORATION UPON OUTGO OF CALCIUM AND MAGNESIUM. Soil Sci. 21: 377–391, illus.

(170) ——— (C)
1926. RECIPROCAL REPRESSION BY CALCIC AND MAGNESIC ADDITIONS IN SURFACE SOIL. Amer. Soc. Agron. Jour. 18: 482–497, illus.

(171) ——— (C)
1926. INFLUENCE OF FORM, SOIL-ZONE, AND FINENESS OF LIME AND MAGNESIA INCORPORATIONS UPON OUTGO OF SULPHATES AND NITRATES. Soil Sci. 22: 21–30.

(172) ——— Shaw, W. M., and Sanders, K. B. (C)
1927. THE INFLUENCE OF LIMING ON THE AVAILABILITY OF SOIL POTASH. LYSIMETER INVESTIGATIONS. Amer. Soc. Agron. Jour. 19: 490–505, illus.

(173) Mooers, C. A., MacIntire, W. H., and Young, J. B. (ABC)
1927. THE RECOVERY OF SOIL NITROGEN UNDER VARIOUS CONDITIONS AS MEASURED BY LYSIMETERS OF DIFFERENT DEPTHS. Tenn. Agr. Expt. Sta. Bul. 138, 30 pp., illus.

(174) MacIntire, W. H. (CD)
1927. THE FATE OF FRACTIONAL INCORPORATIONS OF BURNT LIME IN TWO SOIL ZONES. Soil Sci. 24: 475–485, illus.

(175) ——— and Shaw, W. M. (E)
1928. THE ABSORPTION OF $Ca(OH)_2$–$CaSO_4$ BY SOIL AS A POSSIBLE INDEX TO COLLODIAL ALUMINA AND SILICA. 1st Internatl. Cong. Soil Sci. Washington, D. C. 1927, Comn. 2, Proc. and Papers, v. 2, pp. 279–313, illus.

(176) ——— (AE)
1928. PRESIDENT'S ADDRESS: THE ROLE OF THE LYSIMETER IN SOIL CHEMISTRY RESEARCH. Assoc. Off. Agr. Chem. Jour. 11: 16–26, illus.

(177) ——— and Sanders, K. B. (C)
1929. THE RELATION BETWEEN THE ABSORBED AND THE EXCHANGEABLE CALCIUM AND MAGNESIUM CONTENT OF A SOIL FOUR YEARS AFTER ADDITIONS. Soil Sci. 28: 289–304.

(178) Lyon, T. Lyttleton, and Buckman, Harry O. (CD)
1929. the nature and properties of soils; a college text of edaphology. Rev. ed., 428 pp., illus. New York. *See* Amount of Nutrients Removed from Soil by Drainage, pp. 251–255.

(179) Mooers, C. A., and MacIntire, W. H. (D)
1929. an issue raised in "a new type of lysimeter at the new jersey agricultural experiment station." Science 70: 611–612.

(180) MacIntire, W. H., and Shaw, W. M. (C)
1930. lime-magnesia ratios in dolomitic limestones as influencing solution and soil reactions. Amer. Soc. Agron. Jour. 22: 14–27.

(181) ——— and Sanders, K. B. (AC)
1930. the fixation of the potash of a green manure by liming materials. Soil Sci. 29: 109–117, illus.

(182) ——— Shaw, W. M., and Young, J. B. (C)
1930. the repressive effects of lime and magnesia upon soil and subsoil potash. Jour. Agr. Sci. [England] 20: 499–510.

(183) ——— and Shaw, W. M. (C)
1930. normal magnesium carbonate in comparison with "light" and "heavy" oxides and carbonates in the soil. Amer. Soc. Agron. Jour. 22: 919–934, illus.

(184) ———Shaw, W. M., and Young, J. B. (C)
1930. the oxidation of pyrite and sulfur as influenced by lime and magnesia: a 12-year lysimeter study. Soil Sci. 30: 443–457, illus.

(185) Russell, E. J. (E)
1932. soil conditions and plant growth. Ed. 6, 636 pp., illus. London, New York [etc.] *See* Calcium Oxide and Calcium Carbonate, pp. 233–234.

(186) MacIntire, W. H. (E)
1933. chemistry: soil chemistry. Tenn. Agr. Expt. Sta. Ann. Rpt. (1932) 45: 26–29.

(187) ——— Sanders, K. B., and Shaw, W. M. (BC)
1933. the availability of hydrated lime, limestone, and dolomite of two degrees of fineness, with supplements of red clover hay, as measured by lysimeter leachings. Amer. Soc. Agron. Jour. 25: 285–297, illus.

(188) ——— Mooers, C. A., and Young, J. B. (C)
1933. the effects of additions of certain colorado soils upon the nitrogen balance of a tennessee soil. Soil Sci. 36: 361–368, illus.

(189) ——— Shaw. W. M., Robinson, B., and Sanders, K. B. (C)
1933. the effects of additions of certain colorado soils upon the outgo of bases, chlorides, and sulfates from a tennessee soil. Soil Sci. 36: 435–446.

(190) ——— Shaw, W. M., and Robinson, B. (C)
1934. the distinction between magnesium absorbed and that exchangeable, four years after lysimeter incorporations of oxides and carbonates. Soil Sci. 37: 289–303, illus.

(191) ——— Ellett, W. B., Shaw, W. M., and Hill, H. H. (ABCD)
1934. the conservation of burnt lime, limestone, dolomite and calcium silicate in soil as influenced by methods of incorporation. Tenn. Agr. Expt. Sta. Bul. 152, 52 pp., illus.

(192) ——— Shaw, W. M., and Robinson, B. (BC)
1938. the leaching action of rainwater upon dolomite and limestone separates incorporated with quartz in outdoor lysimeters. Soil Sci. 46: 9–19.

(193) ——— Shaw, W. M., Young, J. B., and Robinson, B. (B)
1938. behavior of calcium, magnesium, and potassium sulfates, as influenced by limestone and by dolomite—a lysimeter study. Soil Sci. 46: 229–247, illus.

TEXAS

(194) Fraps, G. S. (ABC)
1915. losses of moisture and plant food by percolation. Tex. Agr. Expt. Sta. Bul. 171, 51 pp., illus.

(195) American Geophysical Union, Hydrology Committee. (E)
1933. list of agencies in the united states carrying on work in the field covered by the committee on absorption and transpiration, with brief information regarding their work and publications. [united states geological survey, department of interior]. Natl. Res. Council, Amer. Geophys. Union Trans. 14: 356. [Processed.]

(196) Deeter, E. B., and Hopkins, P. L. (B)
1938. progress report of the blackland soil and water conservation experiment station, temple, texas, 1931–1936. methods of investigation: lysimeters; water losses from lysimeters. U. S. Soil Conserv. Serv. ESR–7: 20, 31–32. [Mimeographed.]

VIRGINIA

(197) Ellett, W. B., and Hill, H. H. (A)
1928. investigations in agricultural chemistry: the leaching of plant food through the soil. Va. Agr. Expt. Sta. Ann. Rpt. 1919–27: 26–28.

(198) ——— and Hill, H. H. (ACD)
1929. effect of lime materials on the outgo of sulphur from hagerstown silt loam soil. Jour. Agr. Res. 38: 697–711.

(199) ——— Hill, H. H., and Eheart, J. F. (C)
1931. green manure project. lysimeter investigations. Va. Agr. Expt. Sta. Rpt. 1927–31: 21–24.

(200) Hill, H. H. (ABC)
1934. the effects of mulched and turned rye in the green and mature stages on the liberation of plant nutrients from a silt loam soil. Va. Agr. Expt. Sta. Tech. Bul. 53, 18 pp.

(201) MacIntire, W. H., Ellett, W. B., Shaw, W. M., and Hill, H. H. (ABCD)
1934. the conservation of burnt lime, limestone, dolomite and calcium silicate in soil as influenced by methods of incorporation. Va. Agr. Expt. Sta. Tech. Bul. 54, 52 pp., illus.

(202) Hill, H. H. (ABC)
1937. the liberation of plant nutrients from the soil as affected by alfalfa. Va. Agr. Expt. Sta. Tech. Bul. 60, 19 pp., illus.

(203) Ellett, W. B., and Hill, H. H. (ABCD)
1937. the effects of certain lime materials on the leachings from frederick silt loam soil. Va. Agr. Expt. Sta. Tech. Bul. 61, 19 pp.

WASHINGTON

(204) Martin, J. C. (E)
1938. western society of soil science. [lysimetry]. Science 88: 202–203.

WISCONSIN

(205) Lake States Forest Experiment Station. (AB)
1937. forest litter and gravelly soils—effective conservers of water (la crosse, wis.). U. S. Dept. Agr., Lake States Forest Expt. Sta. Tech. Note 124, 1 p. [Mimeographed.]

(206) Hays, O. E., and Palmer, V. J. (A)
1937. progress report upper mississippi valley soil and water conservation experiment station, la crosse, wisconsin, 1932–1935. introduction. U. S. Soil Conserv. Serv. ESR–1, Illus. [Mimeographed.] *See* footnote, p. 2, and figure 3.

(207) Scholz, H. F. (BD)
1938. trials show woodland increases ground water supply. Wis. Agr. Expt. Sta. Ann. Rpt. (1937) 54 (Bul. 440): 89–90.

MISCELLANEOUS

(208) Cameron, F. K. (E)
1912. THE ROLE OF THE LYSIMETER IN SOIL SOLUTION STUDIES. 8th Internatl. Cong. Appl. Chem., Washington-New York, 1912, Sect. 7 Agr. Chem., Orig. Commun. 15: 49–50.

(209) Wright, R. C. (AC)
1916. GROWING PLANTS IN LARGE CONTAINERS UNDER CONTROL CONDITIONS. Amer. Soc. Agron. Jour. 8: 113–116, illus.

(210) American Society of Agronomy. (A)
1933. STANDARDIZATION OF FIELD EXPERIMENTS: RECOMMENDED STANDARDS FOR LYSIMETER EXPERIMENTS. Amer. Soc. Agron. Jour. 25: 807–810.

GREAT BRITAIN

ENGLAND

(211) Dalton, J. (AB)
1802. EXPERIMENTS AND OBSERVATIONS TO DETERMINE WHETHER THE QUANTITY OF RAIN AND DEW IS EQUAL TO THE QUANTITY OF WATER CARRIED OFF BY THE RIVERS AND RAISED BY EVAPORATION, WITH AN INQUIRY INTO THE ORIGIN OF SPRINGS.—AN ESTIMATE OF THE QUANTITY OF WATER RAISED BY EVAPORATION. Mem. Lit. Phil. Soc. (Manchester, March 1799) 5: 360–372.

(212) ——— (AB)
1819. EVAPORATION. EVAPORATION IN METEOROLOGY, 1796–1798. *In* Rees's Cyclopaedia or Univ. Dictionary of Arts, Sci. and Lit. v. 13. London.

(213) Clutterbuck, J. C. (BE)
1844. OBSERVATIONS ON THE PERIODICAL DRAINAGE AND REPLENISHMENT OF THE SUBTERRANEOUS RESERVOIR IN THE CHALK BASIN OF LONDON Inst. Civ. Engin. Minutes Proc. (session 1842) 2: 155–158. [No reference made to lysimeters.] Discussion by Buckland and J. Dickinson (session 1842), pp. 159–162 [BE]; by J. Dickinson and J. C. Clutterbuck (session 1843), pp. 159–162 [BE].

(214) Parkes, J. (ABD)
1845. ON THE QUANTITY OF RAIN COMPARED WITH THE QUANTITY OF WATER EVAPORATED FROM OR FILTERED THROUGH SOIL; WITH SOME REMARKS ON DRAINAGE. Roy. Agr. Soc. England Jour. (1844) 5: 146–158.

(215) Clutterbuck, J. C. (ABE)
1850. ON THE PERIODICAL ALTERATIONS, AND PROGRESSIVE PERMANENT DEPRESSION, OF THE CHALK WATER LEVEL UNDER LONDON. Inst. Civ. Engin. Minutes Proc. (1849) 9: 151–155. [No reference made to lysimeters.] Discussion by Dickinson, pp. 157–158 [AB]; by Homersham, p. 162 [E].

(216) Barlow, P. W. (ABDE)
1855. ON SOME PECULIAR FEATURES OF THE WATER-BEARING STRATA OF THE LONDON BASIN. Inst. Civ. Engin. Minutes Proc. (1854) 14: 51–52 [AB]. Discussions by Hawkshaw, p. 76 [ABD]; and P. W. Barlow, Dickinson, S. C. Homersham, Evans, and Hawkshaw, pp. 80–84, 84–86, 86–88, illus. [ABE].

(217) Braithwaite, F. (AB)
1861. ON THE RISE AND FALL OF THE RIVER WANDLE; ITS SPRINGS, TRIBUTARIES, AND POLLUTION. Inst. Civ. Engin. Minutes Proc. 20: 191–208. [No reference made to lysimeters]. Discussions by J. Evans, pp. 219–228 [AB], and Hawksley, pp. 230–231 [B].

(218) Denton, J. B. (B)
1862. ON THE DISCHARGE FROM UNDER-DRAINAGE, AND ITS EFFECT ON THE ARTERIAL CHANNELS AND OUTFALLS ON THE COUNTRY. Inst. Civ. Engin. Minutes Proc. (1861) 21: 48–82. [No reference made to lysimeters.] Discussion by Evans, pp. 90–92 [B].

(219) Greaves, C. (ABDE)
1876. ON EVAPORATION AND ON PERCOLATION (1851–1873]. Inst. Civ. Engin. Minutes Proc. 45: 19–47, illus. [AB]. Discussions by Evans, pp. 51–52 [E]; J. H. Gilbert, pp. 52–53, 59–60 [ABD]; Ebermayer, pp. 53–54, 56–58 [ABD]; Homersham, pp. 71, 72 [AB]; R. Field, pp. 86–89 [BD]; and Greaves, pp. 92–94 [AE].

(220) Evans, J. (AB)
1876. on the percolation of the rainfall on absorbent soils. Inst. Civ. Engin. Minutes Proc. 45: 208–216.

(221) Lawes, J. B., and Gilbert, J. H. (AB)
1880. our climate and our wheat crops II. the season of 1878–1879, and the experimental wheat-crops at rothamsted. Roy. Agr. Soc. England Jour. (2) 16: 195–210. (Reprint in Rothamsted Expt. Sta., Harpenden, Mem. Agr. Chem. Physiol. (1880) 5: 25–40, London. 1890.)

(222) ——— Gilbert, J. H., and Warington, R. (ABC)
1881. on the amount and composition of the rain and drainage-waters collected at rothamsted. Roy. Agr. Soc. England Jour. (2) 17: 241–79, illus.

(223) ——— Gilbert, J. H., and Warington, R. (ABCD)
1881. on the amount and composition of the rain and drainage-waters collected at rothamsted part II. the amount and composition of the drainage waters from unmanured fallow land. Roy. Agr. Soc. England Jour. (2) 17: 311–350.

(224) ——— Gilbert, J. H., and Warington, R. (BC)
1882. on the amount and composition of the rain and drainage-waters collected at rothamsted III. the quantity of nitrogen lost by drainage. Roy. Agr. Soc. England Jour. (2) 18: 43–71.

(225) ——— Gilbert, J. H., and Warington, R. (BC)
1882. atmosphare und wasser. ueber die menge und die zusammensetzung des regen-und drainwassers von rothamsted. Biedermann's Zentbl. f. Agr. Chem. 11: 217–226, 654–659.

(226) ——— and Gilbert, J. H. (AD)
1890. effects of the drought of 1870 on some of the experimental crops at rothamsted. results relating to the growth of barley. Rothamsted Expt. Sta., Harpenden, Mem. Agr. Chem. Physiol. (1871) 3: 29–44.

(227) Gilbert, J. H. (ABD)
1890. on rainfall, evaporation and percolation. Rothamsted Expt. Sta., Harpenden, Mem. Agr. Chem. Physiol. (1876) 5, 13 pp.

(228) Lawes, J. B., Gilbert, J. H., and Warington, R. (ABC)
1890. on the amount and composition of the rain and drainage waters collected at rothamsted. Rothamsted Expt. Sta., Harpenden, Mem. Agr. Chem. Physiol. (1882) 5: 1–79, 122–150, illus.

(229) Harrison, J. T. (AB)
1891. on the subterranean water in the chalk-formation of the upper thames, and its relation to the supply of london. Inst. Civ. Engin. Minutes Proc. 105: 2–25. [No reference to lysimeters.] Discussion by J. H. Gilbert, pp. 31–48 [AB]; by J. Hopkinson, pp. 48–52 [B].

(230) Hopkinson, J. (ABD)
1892. water and water-supply, with special reference to the supply of london from the chalk of hertfordshire. Hertfordshire Nat. Hist. Soc. Trans. (1891) 6: 137–146.

(231) Mawley, E. (AB)
1892. the percolation of rain through comparatively light and through comparatively heavy soil. Hertfordshire Nat. Hist. Soc. Trans. (1891) 6: 175–178.

(232) Gilbert, J. H. (AB)
1892. atmosphare und wasser. beobachtungen über regen, sickerwasser und verdunstung in rothamsted. [Translation by F. Seyfert.] Biedermann's Zentbl. f. Agr. Chem. 21: 73–76.

(233) Willis, J. J. (BCE)
1893. rainfall and drainage at rothamsted. Gard. Chron. (3) 14: 392.

(234) Bieler, K. (ABC)
1896. die rothamsteder versuche nach dem stande des jahres 1894 I. beobachtungen über regenfall und drainage. Landw. Jahrb. 25: 199–211, illus.

(235) STORER, F. H. (ABD)
1888–97. AGRICULTURE IN SOME OF ITS RELATIONS WITH CHEMISTRY. Ed. [1], v. 1, New York, 1888; ed. 7, 620 pp. 1897. *See* Soakage of Rainwater, and Drain-Gauges, ed. [1], pp. 61–68 [ABD], and Evaporation from Hot Sand, Drain-Gauges, etc., ed. 7, pp. 81–90 [ABD].

(236) GILBERT, J. H. (AB)
1899. OBSERVATIONS ON RAINFALL, PERCOLATION AND EVAPORATION; WITH RESULTS OBTAINED AT ROTHAMSTED, OVER 20 HARVEST YEARS, 1870–1 TO 1889–90, INCLUSIVE. Rothamsted Expt. Sta., Harpenden, Mem. Agr. Chem. Physiol. 7 Sup.: 1–20.

(237) WARINGTON, R. (ABD)
1900. LECTURES ON THE PHYSICAL PROPERTIES OF THE SOIL. 231 pp., illus. Oxford. *See* Percolation, pp. 89–91 [AB]; Evaporation from Bare Soil, pp. 109–112 [BD]; and Relations of Soil to Water, pp. 121–122 [BD].

(238) SCOTT, R. H. (ABDE)
1900. RESULTS OF PERCOLATION EXPERIMENTS AT ROTHAMSTED, SEPTEMBER 1870–AUGUST 1899. Roy. Met. Soc. [London] Quart. Jour. 26: 139–148, illus. [AB]; Discussion by E. Mawley, J. Hopkinson, and R. H. Curtis, pp. 148–149, 150–151 [BDE].

(239) MILLER, N. H. J. (C)
1902. THE AMOUNTS OF NITROGEN, AS NITRATES, AND CHLORINE IN THE DRAINAGE THROUGH UNCROPPED AND UNMANURED LAND. [London] Chem. Soc. Proc. 18: 89–90.

(240) HALL, A. D. (ABE)
1903–8. THE SOIL. Ed. 1, 286 pp., illus. London. 1903. Ed. 2, 1908, 311 pp., illus. *See* Percolation, ed. 1, pp. 72–76 [BE], and ed. 2, pp. 75–79 [ABE].

(241) ——— (ABC)
1905–19. THE BOOK OF THE ROTHAMSTED EXPERIMENTS. Ed. [1], 294 pp., illus. London. 1905; ed. 2, revised by E. J. Russell, 332 pp., illus. 1919. *See* eds. [1] and 2, Meteorological Observations, p. 15 [A]; Proportion of Rainfall Percolating through Bare Soil, pp. 22–23 [AB]; and The Nitrates in Drainage Waters, pp. 229–230 [BC].

(242) MILLER, N. H. J. (ABC)
1906. THE AMOUNT AND COMPOSITION OF THE DRAINAGE THROUGH UNMANURED AND UNCROPPED LAND, BARNFIELD, ROTHAMSTED. Jour. Agr. Sci. [England] 1 (4): 377–399, illus. [Reprint in Rothamsted Expt. Sta., Harpenden, Mem. Agr. Sci. v. 8 (1902–12). Harpenden, 1914.]

(243) HALL, A. D., and MILLER, N. H. J. (BC)
1906. THE EFFECT OF PLANT GROWTH AND OF MANURES UPON THE RETENTION OF BASES BY THE SOIL. LOSS OF CALCIUM CARBONATE ON UNMANURED LAND. Roy. Soc. [London], Proc., Ser. B, 77: 13–17.

(244) LUEDECKE, C. (BE)
1906. DAS VERHALTNIS ZWEISCHEN DER MENGE DES NIEDERSCHLAGES UND DES SICKERWASSERS NACH ENGLISCHEN VERSUCHEN. Breslau. Landw. Inst. Mitt. 3: 615–646, illus. *See* Tables VI–X.

(245) HILGARD, E. W. (B)
1906. SOILS. 593 pp., illus. New York and London. *See* Lysimeters, p. 227.

(246) RUSSELL, E. J. (B)
1908. NOTE ON AN APPARENT SECULAR CHANGE IN THE ROTHAMSTED DRAIN GAUGES. Jour. Agr. Sci. [England] 2: 29–34, illus.

(247) LATHAM, B. (ABDE)
1909. PERCOLATION, EVAPORATION, AND CONDENSATION. Roy. Met. Soc. [London] Quart. Jour. 35: 189–206, illus. [AB]. Discussions by W. Marriott, p. 207 [BD]; and J. E. Clarke, F. C. Bayard, W. N. Shaw, H. Mellish, and J. Hopkinson, pp. 208–209, 211 [DE].

(248) LYON, T. L., and FIPPIN, E. O. (AB)
1909. THE PRINCIPLES OF SOIL MANAGEMENT. 531 pp., illus. New York. *See* Percolation, Evaporation, pp. 191–197, illus.

(249) FRAPS, G. S. (ABD)
1913. PRINCIPLES OF AGRICULTURAL CHEMISTRY. 493 pp., illus. Easton, Pa. *See* Capillary Action, Losses of Water from Soil, Evaporation, pp. 140–146, illus.

(250) LYON, T. L., FIPPIN, E. O., and BUCKMAN, H. O. (ABD)
1915. SOILS THEIR PROPERTIES AND MANAGEMENT. 764 pp., illus. New York. *See* The Lysimeter Method of Studying Gravitational Water, pp. 239–242, illus.

(251) MOSIER, J. G., and GUSTAFSON, A. F. (B)
1917. SOIL PHYSICS AND MANAGEMENT. 442 pp., illus. Philadelphia and London. *See* Lysimeters or Drain Gauges, pp. 220–221.

(252) RUSSELL, E. J., and RICHARDS, E. H. (BC)
1920. THE WASHING OUT OF NITRATES BY DRAINAGE WATER FROM UNCROPPED AND UNMANURED LAND. Jour. Agr. Sci. [England] 10:22–43, illus.

(253) LYON, T. LYTTLETON, and BUCKMAN, HARRY O. (ABCD)
1923–37. THE NATURE AND PROPERTIES OF SOILS; A COLLEGE TEXT OF EDAPHOLOGY. Ed. [1], 588 pp., illus., 1923; rev. ed. 428 pp., illus., 1929; and ed. 3, 392 pp., illus., 1937. New York. *See* Percolation Losses and their Control, ed. [1], pp. 206–210, illus. [BCD]; Evaporation Losses, ed. [1], pp. 216–218, illus. [B]; Percolation Losses and their Control, rev. ed., pp. 214–218 [BD]; Evaporation Losses, rev. ed., pp. 224–226 [B]; Methods of Studying Drainage Water, Amounts of Nutrients Removed from Soil by Drainage, rev. ed., pp. 249–255, illus. [CD]; Thermal Movement of Soil Water, ed. 3, pp. 126–128 [B]; and Percolation-Methods of Study, and Percolation Losses of Water, ed. 3, pp. 146–156, illus. [ABD].

(254) ROBINSON, G. W., and JONES, J. O. (E)
1927. LOSSES OF ADDED PHOSPHATE BY LEACHING FROM N. WELSH SOILS. Jour. Agr. Sci. [England] 17: 94–95.

(255) BARROWS, H. K. (ABD)
1927. WATER POWER ENGINEERING. 734 pp., illus. New York and London. *See* Measurements of Soil Evaporation, pp. 91–95.

(256) NICHOLSON, H. H., and PANTIN, B. (AC)
1929. THE LEACHING OUT OF AUTUMNAL DRESSINGS OF NITROGENOUS FERTILIZERS. Jour. Agr. Sci. [England] 19: 297–301, illus.

(257) RUSSELL, E. J. (AC)
1932. SOIL CONDITIONS AND PLANT GROWTH. Ed. 6, 636 pp., illus. London, New York, [etc.]. *See* The Effect of Soil Conditions on the Nature and Extent of the Changes in the Nitrogen Compounds, pp. 357–358.

(258) KOSHAL, R. S., and FISHER, R. A. (BE)
1934. THE EFFECTS OF RAINFALL AND TEMPERATURE ON PERCOLATION THROUGH DRAIN GAUGES. Jour. Agr. Sci. [England] 24: 105–135, illus.

(259) ROTHAMSTED EXPERIMENT STATION. (A)
1934. METEOROLOGICAL OBSERVATIONS. RAINFALL AND DRAINAGE. Rothamsted Expt. Sta., Harpenden, Rpt. 1934: 86–87.

(260) KEEN, B. A. (E)
1936. THE CIRCULATION OF WATER IN THE SOIL BETWEEN THE SURFACE AND THE LEVEL OF UNDERGROUND WATER. Assoc. Internatl. d'Hydrol. Sci. VI. Gen. Assembly, Edinburgh, September 1936, 22 (pt. 1): 328–331.

(261) ROTHAMSTED EXPERIMENT STATION. (A)
1937. METEROLOGICAL OBSERVATIONS. RAINFALL AND DRAINAGE. Rothamsted Expt. Sta., Harpenden, Rpt. 1937:87.

(262) PATERSON, J. W. (ABC)
1938. SCIENCE IN AGRICULTURE, A DISCUSSION OF SCIENTIFC PRINCIPLES IN THEIR RELATION TO FARM PRACTICE. 288 pp., illus. London, New York, [etc.] *See* Drainage Losses and Fertility, pp. 160–164.

SCOTLAND

(263) HENDRICK, J. (ABCD)
1921. THE MEASUREMENT OF SOIL DRAINAGE. WITH AN ACCOUNT OF THE CRAIBSTONE DRAIN GAUGES. Highland and Agr. Soc. Scot. Trans. (5) 33: 56–79, illus.

(264) North of Scotland College of Agriculture. (ABC)
1921–22. the lysimeters or draingauges. North of Scot. Col. Agr. Guide Expts. Craibstone 1921: 36–39; 1922: 32–33.
(265) ——— (BC)
1923–37. the lysimeter or draingauge. North of Scot. Col. Agr. Guide Expts. Craibstone, 1923: 33–34; 1924: 32–33; 1925: 30–31; 1926: 25–26; 1927: 26–28; 1928: 25–27; 1929: 26–28; 1930: 28–29; 1931: 27–29; 1932: 26–29; 1933: 26–28; 1934: 31–33; 1935: 57, 63–64; 1936: 54, 61–62; 1937: 53, 60–61.
(266) Hendrick, J. (ACD)
1924. drainage investigations at aberdeen. Scot Jour. Agr. 7: 8–18.
(267) Hendrick, J., and Welsh, H. D. (BC)
1928. the substances removed by the drainage from a scottish soil. 1st Internatl. Cong. Soil Sci., Washington, D. C., 1927, Proc. and Papers, Comn. 2, v. 2, pp. 358–366.
(268) Welsh, H. D. (ABCD)
1929. rainfall, and the drainage from a scottish soil. [Gt. Brit.] Min. Agr. and Fisheries and Scot. Dept. Agr., Agr. and Met. Conf. Rpt. 1928 (4405): 37–45. [Mimeographed.]
(269) Robinson, G. W. (ABC)
1929. lysimeter studies at aber, near bangor. [Gt. Brit.] Min. Agr. and Fisheries and Scot. Dept. Agr., Agr. and Met. Conf. Rpt. 1928 (4405): 47–49. [Mimeographed.]
(270) Lyon, T. L., and Buckman, H. O. (CD)
1929–37. the nature and properties of soils; a college text on edaphology. Rev. ed., 428 pp., illus.; ed. 3, 392 pp., illus. New York. *See* Amounts of Nutrients Removed From Soils by Drainage, pp. 251–255, and Percolation Losses of Nutrients, ed. 3, pp. 151–153.
(271) Hendrick, J. (BC)
1930. a soil balance sheet for a rotation. results obtained from the craibstone drain gauges. Highland and Agr. Soc. Scot. Trans. (5) 42: 1–27.
(272) ——— and Welsh, H. D. (BC)
1932. intensive manuring of grassland. results obtained from the craibstone drain gauges. Highland and Agr. Soc. Scot. Trans. (5) 44: 86–96.
(273) Russell, E. J. (CD)
1932. soil conditions and plant growth. Ed. 6, 636 pp., illus. London, New York [etc.]. *See* The Effects of Leaching; Calcium and Hydrogen Soils; and Podsolising, pp. 250–253 [CD], and Relationship of the Soil Population to Plant Growth, pp. 435–436 [C].
(274) Hendrick, J. (ABCD)
1933. lysimeter studies at aberdeen. Mezogazdasagi Kutatasok 6: 500–509.
(275) ——— (E)
1933. lysimeter kiserlotek aberdeenben. Mezogazdasagi Kutatasok 6: 509.
(276) ——— and Welsh, H. D. (B)
1934. the effect of treatment on the composition of grass: results from the craibstone drain gauges. Highland and Agr. Soc. Scot. Trans. (5) 46: 202–223.
(277) Robertson, I. M., and Stewart, A. B. (CE)
1935. the effect of the lower layers of typical scottish soils on the uptake of nutrients by plants. 3d Internatl. Cong. Soil Sci., Oxford, 1935, Comn. 4. [No reference made to lysimeters.] Discussion by A. B. Stewart, v. 3, p. 225.
(278) Hendrick, J., and Welsh, H. D. (ABCE)
1935. a note on the nature of the suspended matter in drainage waters from the craibstone lysimeter, aberdeen. 3d Internatl. Cong. Soil Sci., Oxford, 1935, Trans. 1, Papers, Comn. 5: 293–295 [ABC]. Discussion by J. S. Joffe and J. Hendrick, v. 3, p. 232 [E].
(279) Comber, Norman M. (CD)
1936. an introduction to the scientific study of the soil. Ed. 3, 206 pp., illus. London. *See* Drainage Water, pp. 97–99.

(280) HENDRICK, J., and WELSH, H. D. (ABC)
1938. FURTHER RESULTS FROM THE CRAIBSTONE DRAIN GAUGES. Highland and Agr. Soc. Scot. Trans. (5) 50: 184–202.
(281) NORTH OF SCOTLAND COLLEGE OF AGRICULTURE. (B)
1938. THE LYSIMETERS OR DRAINGAUGES. North of Scot. Col. Agr. Guide Expts. Craibstone 1938: 52, 59–60.
(282) HENDRICK, J., and WELSH, H. D. (AC)
1938. THE COMPOSITION OF THE SUSPENDED MATTER IN THE DRAINAGE OF THE CRAIBSTONE LYSIMETERS. Internatl. Soc. Soil Sci. Trans. A, Comn. 2 and Alkali Subcomn. 7: 26–30.

ASIA

INDIA

(283) HAYMAN, J. M. (ABC)
1906. AMOUNT AND COMPOSITION OF DRAINAGE WATERS, RAIN, DEW, AND CANAL WATER COLLECTED DURING THE YEARS 1903–1904, 1904–1905, AND 1905–1906. United Provs. (Cawnpore) Agr. Sta. Rpt. 1906: 23–29.
(284) ARNOTT, M. H., and LEATHER, J. W. (AB)
1907. THE CONSTRUCTION OF DRAIN GAUGES AT PUSA. India Dept. Agr. Mem., Chem. Ser. 1: 58–77, illus.
(285) HAYMAN, J. M. (ABC)
1908. EXPERIMENTS UNDER NITROGEN INVESTIGATIONS: DRAINGAUGE—AMOUNT AND COMPOSITION OF DRAINAGE WATERS COLLECTED DURING THE YEAR 1906–1907. United Provs. (Cawnpore) Agr.Sta. Rpt. 1907: 57–62, illus.
(286) BURT, B. C. (ABC)
1908–12. EXPERIMENTS UNDER NITROGEN INVESTIGATIONS. DRAIN GAUGE—AMOUNT AND COMPOSITION OF DRAINAGE WATERS COLLECTED DURING THE YEAR 1907–1908 [AND 1909–1912.] United Provs. (Cawnpore) Agr. Sta. Rpts. 1908: 18–22, illus.; 1911: 20–26, illus.; 1912: 22–26, illus.
(287) LEATHER, J. W. (AB)
1909. DRAINAGE. Agr. Res. Inst. Pusa, Rpt. 1907–9: 40–41, illus.
(288) BURT, B. C., and LEATHER, J. W. (BC)
1909. EXPERIMENTS UNDER NITROGEN INVESTIGATIONS. DRAIN GAUGE—AMOUNT AND COMPOSITION OF DRAINAGE WATERS COLLECTED DURING THE YEAR 1907–1908. United Provs. (Cawnpore) Agr. Sta. Rpt. 1909: 22–26, illus. [1908–9 are dates covering the period.]
(289) LEATHER, J. W. (E)
1910–13. DRAINAGE DATA. Agr. Res. Inst. Pusa, Rpts. 1909–10: 23; 1910–11: 27–28; 1911–12: 23–24.
(290) ——— (BC)
1912. RECORDS OF DRAINAGE IN INDIA: DRAIN-GAUGE RECORDS AT CAWNPORE, 1903–1910, AND AT PUSA, 1906–1910. India Dept. Agr. Mem., Chem. Ser. 2: 63–140, illus.
(291) ——— (E)
1914. SOIL PROBLEMS. Agr. Res. Inst. Pusa, Rpt. 1912–13: 20.
(292) BURT, B. C. (BC)
1914–15. DRAINAGE RECORDS. United Provs. (Cawnpore) Agr. Sta. Rpts. 1913: 27–30, illus.; 1914: 40–42, illus.; 1915: 29–31.
(293) SEN, J. (E)
1916–31. METEOROLOGY AND DRAINGAUGES. Agr. Res. Inst. Pusa, Rpts. 1915–16: 12–13; 1919–20: 31; 1924–25: 22; 1928–29: 30; 1929–30: 33.
(294) BURT, B. C. (BC)
1916. DRAIN-GAUGE CROPS. United Provs. (Cawnpore) Agr. Sta. Rpts. 1916: 32–33.
(295) DAVIS, W. A. (E)
1917. METEOROLOGY AND DRAINGAUGES. Agr. Res. Inst. Pusa, Sci. Rpts. 1916–17: 23.
(296) HARRISON, W. H. (E)
1918–28. METEOROLOGY AND DRAINGAUGES. Agr. Res. Inst. Pusa, Sci. Rpts. 1917–18: 24; 1918–19: 36; 1921–22: 24; 1922–23: 23; 1926–27: 24; 1927–28: 30.

(297) Warth, F. J. (E)
1921. METEOROLOGY AND DRAINGAUGES. Agr. Res. Inst. Pusa, Sci. Rpts. 1920–21: 21.

(298) Mukerji, J. N. (E)
1924–36. METEOROLOGY AND DRAIN GAUGES. Agr. Res. Inst. Pusa, Sci. Rpts. 1923–24: 19; 1925–26: 26; 1931–32: 97; 1932–33: 108; 1933–34: 111.

(299) Batham, H. N. (ABCD)
1926. DRAINAGE WATERS AT CAWNPORE. India Dept. Agr. Mem., Chem. Ser. 8: 127–152, illus.

(300) Joachim, A. W. R. (ABCD)
1928. THE RESULTS OF DRAINAGE AND LEACHING TRIALS AT PERADENIYA DURING 1927. Trop. Agr. [Ceylon] 70: 302–311, illus.

(301) ——— (AC)
1928–29. REPORT OF THE AGRICULTURAL CHEMIST FOR THE YEAR 1927 [AND 1928.] THE LEACHING OF FERTILIZERS FROM CEYLON SOILS UNDER PERADENIYA CONDITIONS. Ceylon Dept. Agr. Tech. Rpts. 1927: 3 [A]; 1928: 3 [C].

(302) ——— (BC)
1929. THE RESULTS OF DRAINAGE AND LEACHING TRIALS AT PERADENIYA DURING 1928. Trop. Agr. [Ceylon] 73: 265–271.

(303) ——— (ABC)
1930. DRAINAGE AND LEACHING TRIALS AT PERADENIYA 1927 TO 1930. Trop. Agr. [Ceylon] 74: 323–330.

(304) ——— (C)
1930. REPORT FOR THE AGRICULTURAL CHEMIST FOR THE YEAR 1929. LEACHINGS AND DRAINAGE TRIALS AT PERADENIYA. Ceylon Dept. Agr. Tech. (Agr.-Chem.) Rpts. 1929: 4.

(305) ——— (E)
1932. DRAINAGE AND LEACHING TRIALS AT PERADENIYA 1927–1930. Internatl. Soc. Soil Sci., Groningen, Netherlands, 1932, Trans., A (A), Comn. 6: 133. (Reprint of Summary and Conclusions from Tropical Agriculture, June 1930).

(306) Keen, B. A. (E)
1932. METEOROLOGY AND DRAIN GAUGES. Agr. Res. Imp. Inst. Pusa, Sci. Rpts. 1930–31: 44.

(307) Nath, B. V. (E)
1936. METEOROLOGY. DRAINGAUGES AND CONTINUOUS DIFFERENTIAL MANURIAL PLOTS. Agr. Res. Imp. Inst. Pusa, Sci. Rpts. 1934–35: 104.

PALESTINE

(308) Winnik, M. (BCE)
1935. TEN YEARS OF LYSIMETER STUDIES ON SOME SOILS OF PALESTINE. 3d Internatl. Cong. Soil. Sci., Oxford, 1935, Trans. 1, Papers Comn. 4: 212–214 [BC]. Discussion by Hendrick and M. Winnik v. 3, pp. 217–218 [E].

FEDERATED MALAY STATES

(309) Belgrave, W. N. C. (AC)
1929. PERCOLATION EXPERIMENTS. I. NITRIFICATION AND THE EFFECT OF COVER PLANTS. Malayan Agr. Jour. 17: 192–205.

JAPAN

(310) Katutani, M., and Hirata, T. (B)
1928. ON THE MELTING SNOW AND ITS WAYS. Natl. Res. Council (Japan), Jap. Astron. and Geophysics, Abs. Jour. 5 (42): 13. [Abstracted from Japanese in Jour. Met. Relating to Forestry and Hydrol. 9: 9–19. 1927.]

(311) Hirata, T. (E)
1929. CONTRIBUTIONS TO THE PROBLEMS OF THE RELATION BETWEEN THE FOREST AND WATER IN JAPAN. COMPARISON OF THE DISCHARGE OF STREAMS UNDER VARIOUS KINDS OF SURFACE CONDITIONS OF THEIR DRAINAGE AREAS. Imp. Forestry Expt. Sta. Meguro Tokyo, Contrib., p. 23.

(312) TAMATE, S., and HIRATA, T. (E)
1933. THE WATER CONTENTS OF SOILS IN AND OUT OF FOREST AND THEIR RELATIONS TO CERTAIN METEOROLOGICAL ELEMENTS. Natl. Res. Council (Japan), Jap. Astron. and Geophysics Abs. Jour. 10 (72): 15–16. [Abstracted from Japanese in Jour. Met. Relating to Forestry and Hydrol. 12: 41–80. 1932.]

(313) OSUGI, S. (ACE)
1935. LYSIMETER EXPERIMENTS WITH BARLEY, TEA AND RICE. 3d Internatl. Cong. Soil Sci., Oxford, 1935, Trans. 1, Papers Comn. 4: 205–207 [AC]. Discussion by Hendrick v. 3, p. 217 [E].

EUROPE

GERMANY AND AUSTRIA

(314) FRAAS, [C.] (C)
1857. VON DEN LÖSUNGEN IM BODEN. Landw. Ver. in Bayern, München, Ergeb. Landw. u. Agr. Chem. Vers. 1: 56–80.

(315) ——— and ZÖLLER, [R.] (ABC)
1859. WEITERE ERGEBNISSE DER VERSUCHE MIT LYSIMETERN (UNTERIRDISCHEN REGENMESSERN). Landw. Ver. in Bayern, München, Ergeb. Landw. u. Agr. Chem. Vers. 2: 60–84.

(316) HOFFMANN, R. (ABC)
1861. DER BODEN. CHEMISCHE UND PHYSIKALISCHE EIGENSCHAFTEN DES BODENS VERSUCHE MIT LYSIMETERN. Jahresber. über die Fortschr. der Agr. Chem. (1859–60) 2: 9–16.

(317) FRAAS, [C.] (B)
1861. FORTSETZUNG DER ANGABEN DER LYSIMETER. Landw. Ver. in Bayern, München, Ergeb. Landw. u. Agr. Chem. Vers. 3: 79–82.

(318) ZÖLLER, R. (BC)
1861. DIE CHEMISCHE UNTERSUCHUNG DER LYSIMETER-RÜCKSTÄNDE. Landw. Ver. in Bayern, München, Ergeb. Landw. u. Agr. Chem. Vers. 3: 82–119.

(319) ——— and FRAAS, [C.] (C)
1861. DIE ANBAUVERSUCHE. Landw. Ver. in Bayern, München, Ergeb. Landw. u. Agr. Chem. Vers. 3: 119–133.

(320) MOLLENDORFF, G. (AB)
1862. DIE REGENVERHÄLTNISSE DEUTSCHLANDS UND DIE ANWENDBARKEIT DER REGENBEOBACHTUNGEN BEI ENT- UND BEWÄSSERUNGEN UND GEWERBLICHEN ANLAGEN. VERSUCHE MIT APPARATEN. Abhandl. Naturf. Gesell. Gorlitz 11: 140–168.

(321) LIEBIG, J. VON. (BC)
1862. DIE CHEMIE IN IHRER ANWENDUNG AUF AGRICULTUR UND PHYSIOLOGIE. DIE NATURGESETZE DES FELDBAUES. Ed. 7, pt. 2, 469 pp., illus. Braunschweig. *See* Der Boden, pp. 94–101 [BC], and Untersuchung von Drain-, Lysimeter-, Fluss- und Moorwasser, pp. 382–391 [C].

(322) HOFFMANN, R. (BC)
1863. CHEMISCHE UND PHYSIKALISCHE EIGENSCHAFTEN DES BODENS. Jahresber. über die Fortschr. der Agr. Chem. (1861–62) 4: 1–21.

(323) EBERMAYER, E. (AB)
1873. DIE PHYSIKALISCHEN EINWIRKUNGEN DES WALDES AUF LUFT UND BODEN UND SEINE KLIMATOLOGISCHE UND HYGIENISCHE BEDEUTUNG. vol. 1, pt. 2, Resultate der Beobachtungen, 266 pp., and Atlas, 253 pp. Berlin. *See* Verhalten des Regen- und Schneewassers zum Boden, oder Einfluss des Waldes und der Streudecke auf die Bodenfeuchtigkeit, pt. 2, pp. 215–234 [AB], and *in* Atlas, pp. 230–242 [B].

(324) GREAVES, C. (ABDE)
1876. ON EVAPORATION AND ON PERCOLATION [1851–1873]. Inst. Civ. Engin. Minutes Proc. 45: 19–47, illus. [AB]. Discussions by J. Evans, pp. 51–52 [E]; J. H. Gilbert, pp. 52–53, 59–60 [ABD]; Ebermayer, pp. 53–54, 56–58 [ABD]; Homersham, pp. 71–72 [AB]; R. Field, pp. 86–89 [BD]; and Greaves, pp. 92–94 [AE].

(325) EBERMAYER, [E]. (AE)
1879. WIE KANN MAN DEN EINFLUSS DER WÄLDER AUF DEN QUELLENREICHTHUM ERMITTELN? Forstwiss. Centbl. (n. s. 1) 23: 77–81.

(326) Wollny, E. (DE)
1881. HYDROMETEORE. BEOBACHTUNGEN ÜBER DIE EINSICKERUNG UND VERDAMPFUNG DES BODENWASSERS SOLLEN IN DAS SYSTEM DER AGRARMETEOROLOGIE EINBEZOGEN WERDEN; DABEI SIND JEDOCH LYSIMETER KEINESWEGS IN ANWENDUNG ZU BRINGEN. Wollny's Forsch. Agr.-Phys. 4: 292–294, 297, 299–301.

(327) Wohltmann, F. (ABCD)
1887. EIN BEITRAG ZUR PRÜFUNG UND VERVOLLKOMMNUNG DER EXACTEN VERSUCHSMETHODE ZUR LÖSUNG SCHWEBENDER PFLANZEN- UND BODENKULTURFRAGEN. Halle Univ., Landw. Inst. Ber., Phys. Lab. u. Versuchsanst. 2 (7): 1–91, illus.

(328) Wollny, E. (AB)
1888. UNTERSUCHUNGEN ÜBER DEN EINFLUSS DER PFLANZENDECKE UND DER BESCHATTUNG AUF DIE PHYSIKALISCHEN EIGENSCHAFTEN DES BODENS. Wollny's Forsch. Agr.-Phys. 10: [261]–344, illus.

(329) ——— (AB)
1888–90. FORSTLICH-METEOROLOGISCHE BEOBACHTUNGEN. DIE FEUCHTIGKEITSVERHÄLTNISSE DER STREUDECKE. Wollny's Forsch. Agr.-Phys. 10: 428–439 [AB]; 13: 134–143 [B].

(330) ——— (ABC)
1888. UNTERSUCHUNGEN ÜBER DIE SICKERWASSERMENGEN IN VERSCHIEDENEN BODENARTEN. Wollny's Forsch. Agr.-Phys. 11: 1–68 [ABC]; 13: 1–15 [AB]; 19: 212–223 [B].

(331) Ebermayer, E. (ABD)
1889. INFLUENCE DE LA FORÊT ET DE LA CONSISTANCE DES PEUPLEMENTS SUR LE DEGRÈ D'HUMIDITÉ DU SOL ET SUR LA QUANTITÉ D'EAU INFILTRÉE. Ann. de la Sci. Agron. Franç. et Etrang. (1889) 1: 424–454.

(332) ——— (AB)
1889. EINFLUSS DES WALDES UND DER BESTANDESDICHTE AUF DIE BODENFEUCHTIGKEIT UND AUF DIE SICKERWASSERMENGE. Allg. Forst. u. Jagd-Ztg. 65: 1–13.

(333) Wollny, E. (B)
1890. UNTERSUCHUNGEN ÜBER DEN EINFLUSS DER STREUDECKE AUF DIE ERWÄRMUNG UND DURCHFEUCHTUNG DES BODENS. DIE SICKERWASSERMENGEN IN DEM STREUBEDECKTEN BODEN. Wollny's Agr.-Phys. 13: 176–184.

(334) ——— (B)
1890. UNTERSUCHUNGEN ÜBER DAS VERHALTEN DER ATMOSPHÄRISCHEN NIEDERSCHLÄGE ZUR PFLANZE UND ZUM BODEN. Wollny's Agr. Phys. 13: 316–356, illus.

(335) Wohltmann. F. (BC)
1891. EIN BEITRAG ZUR PRÜFUNG UND VERVOLLKOMMNUNG DER EXACTEN VERSUCHSMETHODE ZUR LÖSUNG SCHWEBENDER PFLANZEN- UND BODENKULTUR FRAGEN. FORTSETZUNG ENTHALTEND DIE VERSUCHSAUSFÜHRUNG UND RESULTATE DES JAHRES 1886. Halle Univ., Landw. Inst. Ber., (Phys. Lab. und Versuchsanst. 1886) 2 (8): 21–47.

(336) Scheffler, H. (E)
1891. EIN BEITRAG ZUR PRÜFUNG UND VERVOLLKOMMNUNG DER EXACTEN VERSUCHSMETHODE ZUR LÖSUNG SCHWEBENDER PFLANZEN- UND BODENKULTUR FRAGEN. FORTSETZUNG ENTHALTEND DIE VERSUCHSAUSFÜHRUNG UND RESULTATE DES JAHRES 1887. Halle Univ., Landw. Inst. Ber. (Phys. Lab. and Versuchsanst. 1887) 2 (8): 48–69.

(337) ——— (ABCD)
1891. DAS DRAINAGEWASSER UND DIE DURCH DASSELBE HERVORGERUFENEN VERLUSTE AN PFLANZENNÄHRSTOFFEN. Halle Univ., Landw. Inst. Ber. (Phys. Lab. u. Versuchsanst. 1887) 2 (8): 69–120, illus.

(338) Wohltmann, F. (ABC)
1891. EIN BEITRAG ZUR PRÜFUNG UND VERVOLLKOMMNUNG DER EXACTEN VERSUCHSMETHODE ZUR LÖSUNG SCHWEBENDER PFLANZEN- UND BODENKULTUR FRAGEN. FORTSETZUNG. ZUSAMMENSTELLUNG DER ERGEBNISSE DER VERSUCHSJAHRE 1885, 1886, 1887. Halle Univ., Landw. Inst. Ber. (Phys. Lab. und Versuchsanst. 1887) 2 (8): 121–168, illus.

(339) WOLLNY, E. (AB)
1893. UNTERSUCHUNGEN ÜBER DEN EINFLUSS DER MÄCHTIGKEIT DES BODENS AUF DESSEN FEUCHTIGKEITSVERHÄLTNISSE. Wollny's Agr.-Phys. 16: 1–14.

(340) ——— (AB)
1893. UNTERSUCHUNGEN ÜBER DEN EINFLUSS DER STRUKTUR DES BODENS AUF DESSEN FEUCHTIGKEITSVERHÄLTNISSE. Wollny's Agr.-Phys. 16: 381–407.

(341) ——— (AB)
1895. UNTERSUCHUNGEN ÜBER DIE FEUCHTIGKEITSVERHÄLTNISSE DER BODENARTEN. Wollny's Agr.-Phys. 18: 27–62.

(342) ——— (B)
1895. UNTERSUCHUNGEN ÜBER DAS VERHALTEN DER ATMOSPHÄRISCHEN NIEDERSCHLÄGE ZUR PFLANZE UND ZUM BODEN. Wollny's Agr.-Phys. 18: 180–204.

(343) ——— (AB)
1895. UNTERSUCHUNGEN ÜBER DIE VERDUNSTUNG. Wollny's Agr.-Phys. 18: 486–516.

(344) STORER, F. H. (ABD)
1888–97. AGRICULTURE IN SOME OF ITS RELATIONS TO CHEMISTRY. Ed. [1] v. 1, New York 1888; ed. 7, v. 1, 1897. *See* Drain-gauges, ed. [1], pp. 62–68 [BD]; Significance of Natural Mulches in Woodland, ed. [1], pp. 169–175 [AB]; Drain-Gauges. The Bavarian Experiments, ed. 7, pp. 83–87 [ABD]; and Wetness of the Soil in Forests, ed. 7, pp. 274–279 [AB].

(345) WOLLNY, E. (AB)
1897. DIE ZERSETZUNG DER ORGANISCHEN STOFFE UND DIE HUMUSBILDUNGEN MIT RÜCKSICHT AUF DIE BODENCULTUR. 479 pp., illus. Heidelberg. *See* Die Cultur der Humusböden, and Die Moorböden, pp. 368–371 [AB].

(346) HANAMANN, J. (ABC)
1898. LYSIMETER-VERSUCHE. Ztschr. f. das Landw. Versuchsw. Oesterr. 1: 399–410.

(347) TACKE, IMMENDORFF, and MINSSEN. (AC)
1898. UNTERSUCHUNGEN ÜBER DIE ZUSAMMENSETZUNG DER SICKERWÄSSER AUS NICHTGEDÜNGTEM UND AUS GEDÜNGTEM MOORBODEN MIT BESONDERER BERÜCKSICHTIGUNG DER STICKSTOFFVERBINDUNGEN. Landw. Jahrb. 27 (Sup. 4): 349–391.

(348) EBERMAYER, E. (B)
1900. EINFLUSS DER WÄLDER AUF DIE BODENFEUCHTIGKEIT, AUF DAS SICKERWASSER, AUF DAS GRUNDWASSER, UND AUF DIE ERGIEBIGKEIT DER QUELLEN BEGRÜNDET DURCH EXAKTE UNTERSUCHUNGEN. EIN BEITRAG ZU DEN NATURGESETZLICHEN GRUNDLAGEN DES WALDBAUES. 51 pp. Stuttgart. *See* Einfluss der Walder auf die Sickerwässermenge, pp. 29–34.

(349) HANAMANN, J. (BC)
1901. MITTEILUNG DER FÜRSTL. SCHWARZENBERG'SCHEN VERSUCHSSTATION IN LOBOSITZ. LYSIMETERVERSUCHE DES JAHRES 1899. Ztschr. f. das Landw. Versuchsw. Oester. 4: 34–39.

(350) CREYDT, H. O., SEELHORST, [C.] VON, and WILMS. (BC)
1901. UNTERSUCHUNGEN ÜBER DRAINAGE-WASSER. Jour. f. Landw. 49: 251–275.

(351) SEELHORST, C. VON (A)
1902. MITTEILUNGEN VOM LANDWIRTSCHAFTLICHEN VERSUCHFELDE DER UNIVERSITÄT GÖTTINGEN. VEGETATIONSKÄSTEN ZUM STUDIUM DES WASSERHAUSHALTES IM BODEN. Jour. f. Landw. 50: 277–280, illus.

(352) ——— and FRESENIUS. (BC)
1904. BEITRÄGE ZUR LÖSUNG DER FRAGE NACH DEM WASSERHAUSHALT IM BODEN UND NACH DEM WASSERVERBRAUCH DER PFLANZEN. Jour. f. Landw. 52: 355–393.

(353) ——— and MUTHER, [A.] (BC)
1905. MITTEILUNGEN DES LANDWIRTSCHAFTLICHEN VERSUCHSFELDES DER UNIVERSITÄT GÖTTINGEN. I. BEITRÄGE ZUR LÖSUNG DER FRAGE NACH DEM WASSERHAUSHALT IM BODEN UND NACH DEM WASSERVERBRAUCH DER PFLANZEN. Jour. f. Landw. 53: 239–259.

(354) SEELHORST, C. VON (B)
1905. MITTEILUNGEN DES LANDWIRTSCHAFTLICHEN VERSUCHSFELDES DER UNIVERSITÄT GÖTTINGEN. II. UNTERSUCHUNGEN ÜBER DAS EINDRINGEN VON REGENWASSER AUF EINEM SANDBODEN UND AUF EINEM LEHMBODEN. Jour. f. Landw. 53: 260–263.

(355) ——— (B)
1906. MITTEILUNGEN VOM LANDWIRTSCHAFTLICHEN VERSUCHSFELDES DER UNIVERSITÄT GÖTTINGEN. II. ÜBER DEN WASSERVERBRAUCH VON ROGGEN, GERSTE, WEIZEN UND KARTOFFELN. I. MITTEILUNG. Jour. f. Landw. 54: 316–342, illus.

(356) GERLACH. (A)
1908. DAS KAISER WILHELM'S INSTITUT FÜR LANDWIRTSCHAFT IN BROMBERG: LYSIMETERANLAGE. Landw. Jahrb. 37: 198–199. See Tafel IX.

(357) KRÜGER, E. (A)
1910. DIE LYSIMETER DER ABTEILUNG FÜR MELIORATIONSWESEN DES KAISER WILHELM'S INSTITUTS FÜR LANDWIRTSCHAFT IN BROMBERG. Kaiser-Wilhelm-Inst. f. Landw. Mitt. 2: 151–155, illus.

(358) ——— (BC)
1910. LYSIMETERVERSUCHE 1909. Kaiser-Wilhelm-Inst. f. Landw. Mitt. 2: 156–165.

(359) GERLACH. (ABC)
1910. UNTERSUCHUNGEN ÜBER DIE MENGE UND ZUSAMMENSETZUNG DER SICKERWÄSSER. Kaiser-Wilhelm-Inst. f. Landw. Mitt. 2: 319–387, illus.

(360) ——— (ABC)
1910. UEBER DIE DURCH SICKERWÄSSER DEM BODEN ENTZOGENEN MENGEN WASSER UND NÄHRSTOFFE. Illus. Landw. Ztg. 30: [879]–881, illus.

(361) SEELHORST, C. VON. (B)
1910. MITTEILUNGEN AUS DEM LANDWIRTSCHAFLICHEN VERSUCHSFELD DER UNIVERSITÄT GÖTTINGEN. DER WASSERVERBRAUCH VON WIESE UND WEIDE. WASSERVERBRAUCH VON ROGGEN AUF SANDBODEN. Jour. f. Landw. 58: 83–92.

(362) KRÜGER, E. (B)
1911–13. LYSIMETERVERSUCHE 1910, 1911, 1912. Kaiser-Wilhelm-Inst. f. Landw. Mitt. 3: 163–174, 1911; 4: 129–133, 1912; 5: 247–251, 1913.

(363) GERLACH. (BC)
1911. UNTERSUCHUNGEN ÜBER DIE MENGE UND ZUSAMMENSETZUNG DER SICKERWÄSSER. Kaiser-Wilhelm-Inst. f. Landw. Mitt. 3: 351–381.

(364) ——— (BC)
1911. UEBER DIE DURCH SICKERWÄSSER (DRAINAGEWÄSSER) DEM BODEN ENTZOGENEN MENGEN WASSER UND NÄHRSTOFFE. Illus. Landw. Ztg. 31: 755–756.

(365) SEELHORST, C. VON. (ABC)
1913. DER VERBLEIB DES GRÜNDÜNGUNGSSTICKSTOFFS IM SANDBODEN AUF GRUND VON VEGETATIONS VERSUCHEN. Deut. Landw. Gesell. Arb. 241, 147 pp., illus.

(366) RICHTER, G. (B)
1914. LYSIMETERVERSUCHE 1913. Kaiser-Wilhelm-Inst. f. Landw. Mitt. 6: 212–224.

(367) ——— (B)
1914. DER WASSERHAUSHALT DES BODENS. DIE LYSIMETERANLAGE. Kaiser-Wilhelm-Inst. f. Landw. Jahresber. 1913: 50–51.

(368) SIEGERT, and TREIBICH. (B)
1915. ABTEILUNG FÜR MELIORATIONSWESEN. DIE LYSIMETERANLAGE. Kaiser-Wilhelm-Inst. f. Landw. Jahresber. 1914: 43–45.

(369) LYON, T. L., FIPPIN, E. O., and BUCKMAN, H. O. (C)
1915. SOILS, THEIR PROPERTIES AND MANAGEMENT. 764 pp., illus. New York. *See* Drainage records at Bromberg, p. 370.

(370) BRAUN, E. VON, and DADE, H. (BE)
1918. ARBEITSZIELE DER DEUTSCHEN LANDWIRTSCHAFT NACH DEM KRIEGE. 986 pp. Berlin. *See* Die Bodenbewässerung der Wasserbedarf und die Niederschläge, by E. Krüger, pp. 914–917.

(371) GEILMANN, W. (ABC)
1922. DIE AUSWASCHUNG VON NITRAT- UND AMMONIAK-STICKSTOFF AUF SAND UND LEHMBODEN. Jour. f. Landw. 70: 259–266.

(372) LYON, T. LYTTLETON, and BUCKMAN, HARRY O. (ACD)
1923–29. NATURE AND PROPERTIES OF SOILS; A COLLEGE TEXT OF EDAPHOLOGY. Ed. 1, 588 pp., illus; 1923; rev. ed. 428 pp., illus. 1929. New York. *See* Substitution of Bases, ed. 1, pp. 270–272 [AC]; Nature of the Drainage Water from Soils, rev. ed. pp. 246–249 [CD].

(373) FRECKMANN. (B)
1924. JAHRESBERICHT DER PREUSSISCHEN LANDWIRTSCHAFTLICHEN VERSUCHS- UND FORSCHUNGSANSTALTEN ZU LANDSBERG AU DER WARTHE 1923–24. LYSIMETERVERSUCHE. Landw. Jahrb. 60: 153–156.

(374) ——— (AB)
1924. BERICHT ÜBER DIE TÄTIGKEIT DES INSTITUTS FÜR MELIORATIONSWESEN UND MOORKULTUR. DER VERSUCHSGARTEN UND DIE DARIN DURCHGEFÜHRTEN VERSUCHE. Jahresber. der Preuss. Landw. Vers. Forschunganst. Landsberg a. d. Warthe 1920–23: 22–25.

(375) ——— (B)
1924. ARBEITEN AUS DEM VERSUCHSGARTEN UND DEM VERSUCHSFELDE; LYSIMETERVERSUCHE. Jahresber. der Preuss. Landw. Vers. Forschunganst. Landsberg a. d. Warthe 1923–24: 153–156.

(376) ——— JANERT, STAERK, and SIEGERT. (E)
1925. ARBEITEN AUS DEM VERSUCHSGARTEN UND VOM VERSUCHSFELDE. LYSIMETER- UND GRUNDWASSERGEFÄSSVERSUCHE. Jahresber. der Preuss. Landw. Vers. Forschunganst. Landsberg a. d. Warthe 1924–25: 27.

(377) FABIAN, H. (AB)
1926. DIE VERDUNSTUNGSMESSUNGEN AUF DER SCHEITNIGER SCHLEUSENINSEL IN BRESLAU. VERDUNSTUNG VON GRASLAND. Kulturtechniker 29: 396–410, 439, illus.

(378) GERLACH. (ABC)
1926. UNTERSUCHUNGEN ÜBER DIE MENGE UND ZUSAMMENSETZUNG DER SICKERWÄSSER. Landw. Jahrb. 64: 701–733, illus.

(379) FRECKMANN W., JANERT, STAERK, and SIEGERT. (ABC)
1926. BERICHT ÜBER DIE TÄTIGKEIT DES INSTITUTS FÜR MELIORATIONSWESEN UND MOORKULTUR. Landw. Jahrb. 64 (Sup. 1): 34–61, illus.

(380) ——— JANERT, STAERK, and SIEGERT. (E)
1927. ARBEITEN AUS VERSUCHSGARTEN UND VERSUCHSFELD. VERSUCHE ÜBER DEN WASSERBEDARF VERSCHIEDENER GRÄSER. Jahresber. der Preuss. Landw. Vers. Forschungsanst. Landsberg a. d. Warthe 1926–27: 34.

(381) BARROWS, H. K. (BD)
1927. WATER POWER ENGINEERING. 734 pp., illus. New York and London. *See* Measurements of Soil Evaporation, pp. 91–95.

(382) NIKLAS, H., and SCHROPP, W. (A)
1928. DIE LYSIMETERANLAGE DES AGRIKULTURCHEMISCHEN INSTITUTS DER HOCHSCHULE WEIHENSTEPHAN. Fortschr. der. Landw. 3: 1017–1019, illus.

(383) FRECKMANN, BROUWER, STAERK, and SIEGERT. (B)
1928. ARBEITEN AUS VERSUCHSGARTEN UND VERSUCHSFELD. VERSUCHE ÜBER DEN WASSERBEDARF VERSCHIEDENER GRÄSER. Jahresber. der Preuss. Landw. Vers. Forschungsanst. Landsberg a. d. Warthe 1927–28: 39–41.

(384) ———, BROUWER, STAERK, and SIEGERT. (BC)
1928. BERICHT ÜBER DIE TÄTIGKEIT DES INSTITUTS FÜR MELIORATIONSWESEN UND MOORKULTUR. Landw. Jahrb. 68 (Sup. 1): 35–51.

(385) GERLACH. (E)
1928. DIE GROSSE LANDWIRTSCHAFTLICHE WOCHE. DÜNGER-ABTEILUNG. NÄHRSTOFFVERLUSTE DURCH NIEDERSCHLÄGE IN FEUCHTEN UND TROCKENEN JAHREN. Deut. Landw. Gesell. Mitt. 43: 116.

(386) GERLACH. (BC)
1928–29. NAHRSTOFFVERLUSTE DURCH NIEDERSCHLÄGE IN FEUCHTEN UND TROCKENEN JAHREN. Deut. Landw. Gesell. Mitt. 43: 145–147. [Translation read by W. R. Black, Rainfall and Manural Losses in Wet and Dry Years. [Gt. Brit.] Min. Agr. and Fisheries and Scot. Dept. Agr., Agr. and Met. Conf. Rpt., South Kensington, 1928, pp. 51–55.]

(387) ——— (AC)
1928. DIE VERLUSTE AU WERTVOLLEN PFLANZENNÄHRSTOFFEN DURCH DIE ABFLIESSENDEN SICKERWÄSSER. Ver. der Deut. Zuckerindus. Ztschr. 78: 175–186, illus.

(388) KONEKAMP, A., KONIG, SCHWARZ, KALLABIS, and SIEGERT. (E)
1930. LYSIMETERVERSUCHE. UNTERSUCHUNGEN ÜBER DEN EINFLUSS DER DUNGUNG AUF DEN WASSERVERBRAUCH VON FUTTERPFLANZEN. Jahresber. der Preuss. Landw. Vers. Forschungsanst. Landsberg a. d. Warthe (1929–30): 39–40.

(389) ZUNKER, F. (BE)
1930. MENGE DES SICKERWASSERS. *In* Blanck, E., Handbuch der Bodenlehre (Die Physikalische Beschaffenheit des Bodens), v. 6, pp. 195–198. Berlin.

(390) HELBIG, M. (E)
1930. METHODEN ZUR FESTSTELLUNG DER VERDUNSTUNG DES BODENS. *In* Blanck, E., Handbuch der Bodenlehre (Die Physikalische Beschaffenheit des Bodens), v. 6, pp. 246–247. Berlin.

(391) NIKLAS, H., and SCHROPP, W. (A)
1931. AGRICULTURAL RESEARCH INSTITUTES AND AGRICULTURAL COLLEGES VI. THE AGRICULTURAL-CHEMICAL INSTITUTE WEIHENSTEPHAN OF THE TECHNICAL COLLEGE AT MUNICH. Superphosphate 4: 125–131, illus. [In English, French, and German.]

(392) FRECKMANN, W. (E)
1931. DIE REGELUNG DER WASSERVERHLÄTNISSE IM BODEN. *In* Blanck, E., Handbuch der Bodenlehre (Die Physikalische Beschaffenheit des Bodens). v. 9; pp. 2–8. Berlin.

(393) TORNAU, O. (E)
1931. DAS ZIEL DER BODENBEARBEITUNG PHYSIKALISCHE EIGENSCHAFTEN. *In* Blanck, E., Handbuch der Bodenlehre (Die Physikalische Beschaffenheit des Bodens). v. 9, pp. 97–107, illus. Berlin.

(394) POPP, M. (E)
1931. DIE KÜNSTLICHEN DÜNGEMITTEL UND IHRE EINWIRKUNG AUF DEN BODEN. *In* Blanck E., Handbuch der Bodenlehre (Die Massnahmen zur Kultivierung des Bodens). v. 9, pp. 223–226. Berlin.

(395) KRÜGEL, C., DREYSPRING, C., and HEINRICH, F. (DE)
1932. SICKERVERSUCHE MIT SUPERPHOSPHAT UND THOMASMEHL AN 12 BÖDEN. Ztschr. f. Pflanzenernähr. Düngung u. Bodenk. 25A: 195–203, illus.

(396) BARTELS, J. (AB)
1933. VERDUNSTUNG, BODENFEUCHTIGKEIT UND SICKERWASSER UNTER NATÜRLICHEN VERHÄLTNISSEN. Ztschr. f. Forst u. Jagdw. 65: 204–219, illus.

(397) FRIEDRICH, W., and BARTELS, J. (AB)
1935. MESSUNGEN DER VERDUNSTUNG VOM ERDBODEN. Kulturtechniker 38: 234–251, illus.

(398) PFAFF, C. (ACD)
1937. ÜBER LYSIMETER-VERSUCHE. Fortschr. der Landw. Chem. Forsch. 1936, Forschungsdienst Sonderheft 6: 102–114, illus.

(399) LAATSCH, W. (ABE)
1938. DIE DURCHFEUCHTUNGSWERTE DER DEUTSCHEN SANDBODEN. Ernähr. der Pflanze 34: 3–8, illus. [AB]. Abstract in English, Water Relationships in German Sand Soils, and in Spanish, Los Valores de Humedad en los Suelos Arenosos de Alemania, p. 20 [E].

(400) PFAFF, C. (BC)
1938. SICKERWASSER-VERHÄLTNISSE IN LYSIMETERN BEI ZWISCHENFRUCHTBAU. Fortschr. der Landw. Chem. Forsch. 1937, Forschungsdienst Sonderheft 7: 60–66, illus.

BELGIUM

(401) PETERMANN, A. (ABCD)
1889. CONTRIBUTION À LA CHIMIE ET À LA PHYSIOLOGIE DE LA BETTERAVE À SUCRE. [Gembloux] Inst. Agron. de l'Etat Bul. 45, 47 pp. [ABCD]. Review by E. Wollny, Der Einfluss der Vegetation auf die Menge und Zusammensetzung der Drainwässer, Wollny's Agr. Phys. 13: 238–241 [ABCD].

FRANCE

(402) DE LA HIRE, PHILIPPE. (AB)
1720. MÉMOIRES DE MATHEMATIQUE ET DE PHYSIQUE, TIRES DES REGISTRES DE L'ACADEMIE ROYALE DES SCIENCE, DE L'ANNÉE MDCCIII. REMARQUES SUR L'EAU DE LA PLUIE, SUR L'ORIGINE DES FONTAINES; AVEC QUELQUES PARTICULARITÉS SUR LA CONSTRUCTION DES CITERNES. Hist. de l'Acad. Roy. des Sci. Ann. 1703 (2): 56–59.

(403) BERTHELOT. (AB)
1887. SUR LA FIXATION DIRECTE DE L'AZOTE GAZEUX DE L'ATMOSPHÈRE PAR LES TERRES VÉGÉTALES. [Paris] Acad. des Sci. Compt. Rend. 104: 205–209.

(404) ——— (ABC)
1887. RECHERCHES SUR LE DRAINAGE. [Paris] Acad. des Sci. Compt. Rend. 105: 640–646 [ABC]. Translation by Hecht, Untersuchungen über die Drainwässer, Biedermann's Zentbl. f. Agr. Chem. 17: 1–6 [ABC].

(405) DEHÉRAIN, P. P. (ABC)
1890. RECHERCHES SUR L'ÉPUISEMENT DES TERRES ARABLES PAR LA CULTURE SANS ENGRAIS. ÉTUDE DES EAUX DE DRAINAGE. DEUXIÈME MÉMOIRE. ÉTUDE DES EAUX DE DRAINAGE. Ann. Agron. 16: 337–358 [ABC]. Abstract by E. Wollny, Studien über die Drainwässer, Wollny's Agr. Phys. 13: 402–405 [BC].

(406) ——— (BC)
1891. SUR LA COMPOSITION DES EAUX DE DRAINAGE DES TERRES NUES ET CULTIVÉES. Ann. Agron. 17: 49–82.

(407) ——— (BC)
1892. CONTRIBUTION À L'ÉTUDE DES EAUX DE DRAINAGE; TERRES SANS VÉGÉTATION. Ann. Agron. 18: 273–299.

(408) ——— (ABC)
1893. LES EAUX DE DRAINAGE DES TERRES CULTIVÉES. Ann. Agron. 19: 65–89, illus. [ABC]. Abstract by E. Wollny, Die Drainwässer bebauter Böden, Wollny's Agr. Phys. 16: 230–232 [ABC].

(409) ——— (BC)
1894. LES EAUX DE DRAINAGE DES TERRES CULTIVÉES. EAUX DE DRAINAGE D'HIVER. Ann. Agron. 20: 21–42.

(410) ——— (BC)
1894. LES EAUX DE DRAINAGE DES TERRES CULTIVÉES. LES CASES DE VÉGÉTATION DE MARS 1893 À MARS 1894. Ann. Agron. 20: 449–466.

(411) WILEY, H. W. (AE)
1894. PRINCIPLES AND PRACTICE OF AGRICULTURAL ANALYSIS. I. SOILS. 606 pp., illus. Easton. *See* Lysimetry [E] ed. 1, p. 158; ed. 2, p. 171; ed. 3, p. 186; and Improved Method of Deherain [AE] ed. 1, pp. 167–169, illus.; ed. 2, pp. 178–180, illus.; ed. 3, pp. 194–196, illus.

(412) DEMOUSSY, E. (ABC)
1895. INVESTIGATIONS AT THE GRIGNON AGRICULTURAL EXPERIMENT STATION. EXPERIMENTS ON BARE SOILS. U. S. Off. Expt. Stas. Expt. Sta. Rec. (1893–94) 5: 10–27.

(413) DEHÉRAIN, P. P. (BC)
1895. RECHERCHES SUR LES EAUX DE DRAINAGE DES TERRES CULTIVÉES, MARS 1894–MARS 1895. Ann. Agron. 21: 193–207.

(414) ——— (BC)
1896. LA JACHÈRE. Ann. Agron. 22: 257–263.

(415) ——— (B)
1897. OBSERVATIONS RECUEILLES AUX CASES DE VÉGÉTATION DE GRIGNON. (2ND MEM.). Ann. Agron. 23: 225–229.

(416) ——— (BC)
1897. RECHERCHES SUR LES EAUX DE DRAINAGE DES TERRES NUES ET DES TERRES CULTIVÉES (5TH MEM). Ann. Agron. 23: 241–267.

(417) Dehérain P. P. (B)
1898. le travail du sol. pénétration, emmagasinement et mouvement de l'eau dans le sol. Ann. Agron. 24: 449–481, illus.

(418) ——— (ABC)
1902. traite de chimie agricole. Ed. 1902, 969 pp., illus. Paris. *See* Drainage des Terres Nues et Cultivées, Étudié aux Cases de Végétation de Grignon, pp. 338–340, 584–594, illus.

(419) Demolon, A., and Bastisse, E. (BC)
1933. observations en cases lysimétriques (versailles). [France] Min. de l'Agr., Inst. des Rech. Agron. Rap. 1932: 56–63.

(420) ——— and Bastisse, E.
1934. observations en cases lysimétriques (versailles). [France] Min. de l'Agr. Inst. des Rech. Agron. Rap. 1933: 24–30.

(421) ——— and Bastisse, E. (C)
1934. contribution a l'étude de la mécanique chimique des anions dans le sol. Ann. Agron. (n. s.) 4: 53–76, illus.

(422) [France] Ministère de l'Agriculture. (A)
1934. le centre national de recherches agronomiques de versailles; station centrale d'agronomie (sols et biochimie végétale). [France] Min. de l'Agr. Inst. des Rech. Agron. 1934: 3–4, illus.

(423) Demolon, A., and Bastisse, E. (BC)
1936. observations en cases lysimétriques. *In* Recherches sur la Fertilisation effectuées en 1934 par les Stations Agronomiques [France], pp. 10–17.

(424) ——— (CD)
1935. enseignements tirés d'observations continues en cases lysimétriques. Acad. d'Agr. de France Compt. Rend. 21: 62–68.

(425) ——— and Bastisse, E. (AC)
1936. genèse des colloïdes argileux dans l'altération du granite en cases lysimétriques. Ann. Agron. (n. s.) 6: 803–810.

(426) ——— and Bastisse, E. (BC)
1936. études générales sur la fertilisation. observations météorologiques et lysimétriques. *In* Recherches sur la Fertilisation Effectuées en 1935 par les Stations Agronomiques [France], pp. 1–8.

(427) ——— and Bastisse, E. (C)
1937. observations en cases lysimétriques sur la mobilisation des réserves azotées et minérales des sols. [Paris] Acad. des Sci. Compt. Rend. 204: 1495–1497.

(428) ——— (B)
1937. le climat du sol. Ann. Agron. (n. s.) 7: 625–640, illus.

(429) ——— and Bastisse, E. (BC)
1937. études générales sur la fertilisation. observations en cases lysimétriques. *In* Recherches sur la Fertilisation Effectuées en 1936 par les Stations Agronomiques [France] 10: 12–19.

(430) ——— and Bastisse, E. (C)
1938. sur quelques complexes colloidaux du fer et de l'aluminium ii. rôle des complexes colloïdaux du fer et de l'aluminium dans les processus pédogéniques. Ann. Agron. (n. s.) 8: 25.

(431) ——— and Bastisse, E. (ABCD)
1938. results of lysimeter experiments. [versailles]. Soil Sci. 46: 1–7.

(432) ——— and Bastisse, E. (BC)
1938. études générales sur la fertilisation. observations en cases lysimétriques. *In* Recherches sur la Fertilisation Effectuées en 1937 par les Stations Agronomiques [France] 11: 12–20.

SWITZERLAND

(433) Bühler, A. (AB)
1891–95. untersuchungen über sickerwassermengen. Schweiz. Centralanst. f. das Forstl. Versuchsw. Mitt. 1: 291–332, 1891 [AB]; 4: 203–248, 1895 [B].

(434) RISLER, E., and WÉRY, G. (D)
1903. CONTRIBUTIONS A L'ÉTUDE DU DRAINAGE ET DE L'IRRIGATION. CONSOMMATION D'EAU FAITE PER LES PLANTES. [France] Min. de l'Agr., Inst. Natl. Agron. Ann. 2 (2): 5–9.

(435) ——— and WERY, G. (BD)
1904. IRRIGATION ET DRAINAGES. 516 pp., illus. Paris. *See* Transpiration des Plantes et Quantités d'Eau Apportees par les Pluies, pp. 18–31.

(436) GRÜNIGEN, F. V. (CD)
1938. UEBER DIE NÄHRSTOFFVERLUSTE EINES BODENS BEI FORTGESETZTER VERGLEICHENDER MINERALISCHER UND GÜLLEN-DÜNGUNG. Landw. Jahrb. der Schweiz 52: 663–664.

ITALY

(437) DRAGHETTI, A. (E)
1930. RELAZIONE TECNICA SULL'ORGANIZZAZIONE E SUL FUNZIONAMENTO DELLA R. STAZIONE AGRARIA SPERIMENTALE DI MODENA; ATTIVITÀ DELLA STAZIONE AGRARIA. Modena R. Staz. Sper. Agr. Ann. Bol. delle Ricerche e della Sper. (1927–29) 1: 17–19, illus.

(438) ——— (ABC)
1932. RICERCHE SULLA CONCIMAZIONE AZOTATA DIRETTA AL FRUMENTO IN CASSE DA VEGETAZIONE LISIMETRICHE. Modena R. Staz. Sper. Agr. Ann. Bol. delle Ricerche e della Sper. (1930–31) 2: 164–241, illus.

(439) ——— (A)
1936. CICLO DI RICERCHE SULLA SUB-IRRIGAZIONE DELLE PIANTE SARCHIATE E. FORAGGIERE, IN CASSE DA VEGETAZIONE LISIMETRICHE. Modena R. Staz. Sper. Agr. Ann. Bol. delle Ricerche e della Sper. 5: 407–414, illus.

(440) ——— and PANTOLI, B. (AB)
1936. RICERCHE SULLA SUB-IRRIGAZIONE DEL GRANTURCO IN CASSE DA VEGETAZIONE LISIMETRICHE. Modena R. Staz. Sper. Agr. Ann. Bol. delle Ricerche e della Sper. 5: 415–434.

(441) GIANDOTTI, M. (E)
1937. TEMPERATURA ED EVAPORAZIONE DEL SUOLO E DELLE ACQUE. EVAPORAZIONE DEL SUOLO. Idrologia della Bonifica Integrale, Biblioteca della Bonifica Integrale 3 (pt. 1): 79–80.

(442) GIGLIOLI, G. R. (E)
1938. L'EROSIONE DEL TERRENO AGRARIO NEI TROPICI. IL PROBLEMA DELLA DECADENZA DELLA FERTILITÁ NEI TERRENI TROPICALI L'EROSIONE. Inst. Agr. Colon. Ital. Relaz. e. Monog. Agr. Colon. 49: 24. [Views of Thompson's and Musgraves's lysimeters.]

NETHERLANDS

(443) STERK, A. E. (AB)
1898. OVER REGEN VERDAMPING EN KWEL IN DEN HAARLEMMERMEERPOLDER. ONDERGRONDSCHE CONDENSATIE. K. Inst. van Ingen. Tijdschr. Verhandel. 1897–98: 65–67, 74–75.

(444) BRUIJN, H. E. DE. (AB)
1903. OVER LYSIMETER-WAARNEMINGEN EN DE HOEVEELHEID DRINKWATER, DIE DE DUINEN DIENOVEREENKOMSTIG KUNNEN GEVEN. Nederland. Natur- en Geneesk. Cong. Handel. 9: 147–154, illus.

(445) DEUTSCHE LANDWIRTSCHAFTS-GESELLSCHAFT. (ABCD)
1906. LANDWIRTSCHAFTLICHE ERFAHRUNGEN DES AUSLANDES. BELEUCHTUNG DER BODENNITRIFIKATION DURCH DRAINWASSERUNTERSUCHUNGEN. 35 JÄHRIGE LYSIMETERVERSUCHE IN ROTHAMSTED UND NEUERE UNTERSUCHUNGEN IN HOLLAND. Deut. Landw. Gesell. Mitt. 21: 195–203, illus.

(446) HUDIG, J. (ABCD)
1906. NITRIFICATIE EN DE SAMENSTELLING VAN DRAINWATER. Cultura 18: 149–163, illus.

(447) VERSLUYS, J. (ABD)
1916. DE CAPILLAIRE WERKINGEN IN DEN BODEM. 136 pp., illus. Amsterdam. *See* Lysimeterwaarnemingen, pp. 116–129 [ABD].

(448) Hesselink, E., and Hudig, J. (AB)
1925. de invloed van eene bodembedekking bij stuifzand op den groei der dennen. Rijksboschbouwproefsta. Wageningen, Meded. 2(2): 129–172, illus.

(449) ——— and Hudig, J. (E)
1925. vergelijking van de resultaten der verschillende proeven, voor zooveel betreft de lengtegroei. Rijksboschbouwproefsta. Wageningen, Meded. 2 (2): 173–177.

(450) ——— and Hudig, J. (AB)
1925. der einfluss einer bodenbedeckung auf das wachstum von kiefern im flugsand. Rijksboschbouwproefsta. Wageningen, Meded. 2 (2): 178–184.

(451) ——— and Hudig, J. (AE)
1925. die kondensation von atmosferischer feuchtigkeit im bodem. Rijksboschbouwproefsta. Wageningen, Meded. 2 (2): 185–190, illus.

(452) Maschhaupt, J. G. (ABCD)
1938. lysimeter-onderzoekingen aan het rijkslandbouwproefstation te groningen en elders i. regenval, drainage en verdamping. Netherlands Landbouw. Directie van den Verslagen van Landbouwkundige Onderzoekingen. Groningen Rijklandb. Proefsta. 44 (1) A: 1–169, illus., [ABCD]. English summary, Lysimeter Investigations at Groningen and Elsewhere, Part 1, Rainfall, Drainage and Evaporation, pp. 170–179 [ABD].

SCANDINAVIA

(453) Feilitzen, H. von. (ABC)
1900. försöksverksamheten vid rothamsted i england. iakttagelser öfver nederbörd och dränering. Svenska Mosskulturför. Tidskr. [14]: 28–31, illus.

(454) ——— (A)
1905. försöksgården for växtkultur i jönköping och planen till vegetations-försöken år 1905. lysimetrarne. Svenska Mosskulturför. Tidskr. 19: 401–423, illus.

(455) ——— (A)
1910. svenska mosskulturföreningens institutionsbyggnad. vegetationsgarden. Svenska Mosskulturför. Tidskr. 24: 179–184, illus.

(456) ——— Lugner, I., and Hjerstedt, H. (?)
1912. einige untersuchungen über die mit den sickerwassern aus umbebautem und mit verschiedenen kulturpflanzen bebautem moorboden entstandenen verluste an pflanzennahrstoffen. Kulturtechniker 1: 210–220.

(457) ——— Lugner, I., and Hjerstedt, H. (ABCD)
1912. några undersökningar öfver de mängder växtnäringsämnen, som med nederbörden uttvättas och gå förlorade från obevuxen och med olika kulturväxter bevuxen torfjord. resultat af 5-åriga lysimeter-försök i vegetationsgården i jönköping. Svenska Mosskulturför. Tidskr. 26: 111–154, illus.

(458) ——— (AB)
1913. svenska mosskulturföreningens kulturförsök i jönköping, vid flahult och torestorpsmossen år 1911; hvilken inverkan utöfvar ett olika högt grundvattenstånd på gräsvallars utveckling och gräsbeståndets sammansättning på torfjord? Svenska Mosskulturför. Tidskr. 27: 3–8, illus.

SOVIET UNION

(459) Malüschitzki, N. (AB)
1900. k voprosu o znachenii. evaporometricheskiskh pokazanii dlia zaprosocselsko-khoziastvennoi. praktiki 3, nabludenia nad espareneiamy vlagy pochvoю. Ann. Inst. Agron. de Moscu. 6: 349–368, illus.

(460) Welbel, B. (BC)
1901. otchet po khimicheskoĭ laboratoriy. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1900) 6: 77–103. [Tabular data in Russian and French.]

(461) [TROUBETZKOY, L.] (C)
1901. EXTRAIT DU COMPTE RENDU DU LABORATOIRE CHIMIQUE POUR L'ANÉE 1900; LES EAUX LYSIMÉTRIQUES. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1900) 6: 223–231.

(462) KOSSOVICH, P., and TRET'IAKOV, I. (AB)
1902. KVOPROSU O VLIIANII UGLEKISLAGO KAL'TSIIA NA BYSTROTU RAZLOJENIÎA ORGANICHESKIKH VESHCHESTV. Zhur. Opytn. Agron. (Jour. f. Expt. Landw.) 3: 450–476 [AB]. Review by Kossovich, P., Zur Frage über den Einfluss des kohlensauren Calciums auf den Gang der Zersetzung organischer Stoffe, pp. 476–484 [AB]

(463) WELBEL, B. (ABC)
1903. DIE LYSIMETERWASSER. Review by L. Richter. Biedermann's Zentbl. f. Agr. Chem. 32: 293–296.

(464) ——— (BC)
1903. OTCHET PO KHIMICHESKOÏ LABORATORIY. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1902) 8: 69–96, [Tabular data in Russian and French.]

(465) WINKLER, A. (BC)
1903. RESUMÉ DU COMPTE RENDU DU LABORATOIRE DE CHIMIE POUR L'ANNÉE 1902. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1902) 8: 143–151.

(466) WELBEL, B. (BC)
1903. EZOOCHENEIA SOSTOVA LEZOMETRICHESKIKH VOD E MEMERALEZATSIA POCHVENAVO AZOTA. Zhur. Opytn. Agron. (Jour. f. Expt. Landw.) (1903) 4: 285–307. [Tabular data in Russian and German.]

(467) ——— (C)
1903. BEITRÄGE ZUM STUDIUM DES LYSIMETERWASSERS UND DER NITRIFICATION DES BODENSTICKSTOFFS. Zhur. Opytn. Agron. (Jour. f. Expt. Landw.) (1903) 4: 307–312.

(468) ——— (B)
1904. IZSLIEDOVANEIA KHIMICHESKOI LABORATORIY. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1903) 9: 95–101. [Tabular data in Russian and French.]

(469) WINKLER, A. (C)
1904. RESUMÉ DU COMPTE RENDU DU LABORATOIRE CHIMIQUE POUR L'ANNÉE 1903. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1903) 9: 130–134.

(470) WELBEL, B. (BC)
1905. IZSLIEDOVANEIA KHIMICHESKOÏ LABORATORIY. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1904) 10: 75–116. [Tabular data in Russian and French.]

(471) WINKLER, A. (C)
1905. RÉSUMÉ DU COMPTE RENDU DU LABORATOIRE DE CHIMIQUE POUR L'ANNÉE 1904. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1904) 10: 147–160.

(472) WELBEL, B. (BC)
1906. ATMOPHERNIY OSADKY. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1905) 11: 73–119, illus. [Tabular data in Russian and French.]

(473) WINKLER, A. (C)
1906. RÉSUMÉ DU COMPTE RENDU DU LABORATOIRE DE CHIMIE. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1905) 11: 128–135.

(474) WELBEL, B. M. (ABC)
1906. DANNYIA LEZOMETRICHESKIKH ETSLEDOVANY PLOTIANSKOÏ C-KHOZ. OPUITNOY STANTZII. Zhur. Opytn. Agron. (Jour. f. Expt. Landw.) (1905) 6: 129–163 [ABC]. [Tabular data in Russian and German], and review, Daten Lysimetrischer Untersuchungen von der Versuchsstation Ploty, pp. 163–175 [ABC].

(475) TROUBETSKOY, L., and BYTCHIKHINE, A. (ABC)
1906. POSITION GÉOGRAPHIQUE DE LA STATION AGRONOMIQUE ET SON ORGANISATION; L'ÉVAPORATION ET LA FILTRATION DE L'EAU DANS LE SOL; ACTIVITÉ NITRIFIANTE DU SOL. Ann. de la Sci. Agron. (3) 1: 332–336, 342–346, 389–395.

(476) Welbel, B. (C)
1907. lysimetrische untersuchungen an der versuchstation ploty. Review by L. Richter. Biedermann's Zentbl. f. Agr. Chem. 36: 782.

(477) ——— (BC)
1907. izsliedovaneia khimicheskoĭ laboratoriy opuitn. stantzii. process at nytriphikatziy e denitriphikatzi v pochva po lezometrichesky izsliedovaniam. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1906) 12: 148–208.

(478) Winkler, A. (C)
1907. résumé du compte-rendu du laboratoire de chimie. Ploty Selsk. Khoz. Opytn. Sta. (Ploty. Sta. Expt. Agron. Ann. Rap.) (1906) 12: 248–262.

(479) Welbel, B. (BC)
1908. process nytriphikatziy v pochva yarovogo, parovogo e ozimogo kleena. Ploty Selsk. Khoz. Opytn. Sta. (Polty Sta. Expt. Agron. Ann. Rap.) (1907) 13: 105–146.

(480) Winkler, A. (C)
1908. résumé des travaux du laboratoire, ainsi que des expériences sur la végétation en vases dans les, en 1907. Ploty Selsk. Khoz. Opytn. Sta. (Ploty Sta. Expt. Agron. Ann. Rap.) (1907) 13: 175–191.

(481) Gemmerling, W. W. (AB)
1922. ovodnom regime podzolistihk pochvy na osnovaniy dannykh lezometrichesky izsliedovany. Moskov. Oblastn. Selsk. Khoz. Opytn. Sta. Trudy (Moscow Regional Agr. Expt. Sta. Works) Bul. 1: 88–95, illus. [AB]; also abstract, About the Water Regime of Ashy Loam Soils, According to Lysimeter Investigations, p. 124–125 [A].

(482) Popov, V. P. (AB)
1928. pochvennaîa vlaga e metody chk ezoochenia. Mleev Sadovo-Ogorod. Opytn. Sta. Trudy (Mleev Hort. Expt. Sta. Sect. Rural Met. Bul.) 16: 39–69, illus.

(483) Rutkovsky, V. J. (AE)
1936. study of water circulation in layer between the surface of soil and the ground-water level; lysimetric funnels. Internatl. Hydrol. Sci. Assoc., 6th Gen. Assem. Edimbourg, September 1936, 22: 370–375.

AFRICA

NORTH AFRICA

(484) Yankovitch, L. (AC)
1933. contribution à la meilleure compréhension du problème de l'azote dans les terres nord-africaines. Ann. du Serv. Bot. et Agron. [Tunis] 10: 25–35.

(485) Boeuf, F. (A)
1933. organisation des recherches agronomiques au service botanique et agronomique de tunisie. Ann. du Serv. Bot. et Agron. [Tunis] 10: 123–126, illus.

(486) ——— and Novikoff, V. (ABC)
1935–6. cultures en cases de végétation et en cases lysimétriques au service botanique et agronomique de tunisie. Ann. du Serv. Bot. et Agron. [Tunis] 12–13: [677]–696, illus. *Also in* Acad. d'Agr. de France Compt. Rend. 22 (1936): 463–473, illus.

(487) Miege, E. (ABC)
1937. étude du bilan de l'eau et des elements fertilisants dans quelques sols types du maroc. Ann. Agron. 7: 370–388, illus.

SOUTH AFRICA

(488) Thompson, W. R. (ABCD)
1936. moisture farming in south africa. 260 pp. illus. [Johannesburg]. *See* The Role of Percolation in the Dissipation of Moisture, pp. 192–215, illus.

(489) Haylett, D. G., Ross, J. C., and Dawson, E. S. (AB)
1937. percolation experiments. Univ. Pretoria Soil Erosion and Grassland Experiments, Prog. Rpt. G, 1284: 10–11. [Mimeographed.]

AUTHOR INDEX

[Figures refer to entries in the Bibliography]

Adams, J. E., 148, 149.
American Geophysical Union, Hydrology Committee, 7, 195.
American Society of Agronomy, 210.
Anderson, P. J., 11.
Anonymous, 82.
Arnott, M. H., 284.
Ayres, Q. C., 66.
Babcock, S. M., 95, 96.
Bailey, E. M., 24.
Barker, J. F., 108.
Barlow, P. W., 216.
Barnette, R. M., 41, 42, 43, 44.
Barrows, H. K., 119, 255, 381.
Bartel, F. O., 137.
Bartels, J., 396, 397.
Bastisse, E., 419, 420, 421, 423, 425, 426, 427, 429, 430, 431, 432.
Batham, H. N., 299.
Bayard, F. C., 247.
Beattie, H. G., 128.
Belgrave, W. N. C., 309.
Benson, N., 44.
Berthelot, 403, 404.
Bieler, K., 234.
Bizzell, J. A., 109, 112, 114, 118, 120, 123, 127, 133.
Black, W. R., 386.
Blair, A. W., 25, 26, 27, 28.
Blythe, S. O., 8.
Boeuf, F., 485, 486.
Bortner, C. E., 71.
Braithwaite, F., 217.
Braun, E. von, 370.
Brouwer, 383, 384.
Bruijn, H. E. de, 444.
Buckland, 213.
Buckman, H. O., 106, 115, 178, 250, 253, 270, 369, 372.
Bühler, A., 433.
Burgess, P. S., 1.
Burt, B. C., 286, 288, 292, 294.
Bytchikhine, A., 475.
California Forest and Range Experiment Station, 10.
Cameron, F. K., 208.
Clarke, J. E., 247.
Clutterbuck, J. C., 213, 215.
Collison, R. C., 122, 125, 128, 131.
Collison, S. E., 29, 30, 31, 32, 33.
Comber, N. M., 279.
Connecticut Agricultural Experiment Station, 12, 13, 15, 16, 17, 18, 20, 22.
Creydt, H. O., 350.
Curtis, R. H., 238.
Dade, H., 370.
Dalton, J., 211, 212.
Davis, W. A., 295.
Dawson, E. S., 489.
Dean, H. K., 141, 142, 143, 144, 145.
Deeter, E. B., 196.
Deherain, P. P., 405, 406, 407, 408, 409, 410, 413, 414, 415, 416, 417, 418.
De la Hire, P., 402.
Demolon, A., 419, 420, 421, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432.
DeMoussy, E., 412.
Denton, J. B., 218.
Deutsche Landwirtschafts Gesellschaft 445.
Dickinson, J., 213, 215, 216.
Draghetti, A., 437, 348, 439, 440.
Dreyspring, C., 395.
Duley, F. L., 86, 87.
Ebermayer, E., 219, 323, 324, 325, 331, 332, 348.
Eckart, C. F., 45, 46, 50, 51, 53, 54.
Edwards, E. E., 84.
Eheart, J. F., 199.
Ellett, W. B., 191, 197, 198, 199, 201, 203.
Emery, F. E., 101.
Evans, J., 216, 217, 218, 219, 220, 324.
Fabian, H., 377.
Feilitzen, H. von, 453, 454, 455, 456, 457, 458.
Field, R., 219, 324.
Fippin, E. O., 106, 248, 250, 369.
Fisher, R. A., 258.
Fraas, C., 314, 315, 317, 319.
[France] Ministere de l'Agriculture, 422.
Fraps, G. S., 194, 249.
Freckmann, W., 373, 374, 375, 376 379, 380, 383, 384, 392.
Fresenius, 352.
Friedrich, W., 397.
Garstka, W. U., 138, 139.
Geilmann, W., 371.
Gemmerling, W. W., 481.
Gerlach, 356, 359, 360, 363, 364, 378, 385, 386, 387.
Giandotti, M., 441.
Giglioli, G. R., 442.
Gilbert, J. H., 219, 221, 222, 223, 224, 225, 226, 227, 228, 229, 232, 236, 324.
Goessmann, C. A., 80.
Goff, E. S., 97, 98, 99, 101.
Goss, A., 61.
Greaves, C., 219, 324.
Grunigen, F. V., 436.
Gustafson, A. F., 251.

Hall, A. D., 240, 241, 243.
Hanamann, J., 346, 349.
Harlan, J. D., 128.
Harrison, J. T., 229.
Harrison, W. H., 296.
Hawkshaw, J., 216.
Hawksley, T., 217.
Haylett, D. G., 489.
Hayman, J. M., 283, 285.
Hays, O. E., 206.
Hecht, A., 404.
Hedrick, U. P., 111.
Heinrich, F., 395.
Helbig, M., 390.
Hendrick, J., 263, 266, 267, 271, 272, 274, 275, 276, 278, 280, 282, 308 313.
Hendrickson, H. C., 147.
Hesselink, E., 448, 449, 450, 451.
Hester, J. B., 43.
Higby, W. M., 146.
Hilgard, E. W., 245.
Hill, H. H., 191, 197, 198, 199, 200, 201, 202, 203.
Hirata, T., 310, 311, 312.
Hjerstedt, H., 456, 457.
Hoffmann, R., 316, 322.
Holding, W. A., 154.
Homersham, S. C., 215, 216, 219, 324.
Hopkins, P. L., 196.
Hopkinson, J., 229, 230, 238, 247.
Horton, R. E., 103.
Hudig, J., 446, 448, 449, 450, 451.
Huston, H. A., 61.
Immendorff, 347.
Indiana Agricultural Experiment Station, 60.
Jacobson, H. G. M., 14.
Janert, 376, 379, 380.
Joachim, A. W. R., 300, 301, 302, 303, 304, 305.
Joffe, J. S., 88, 89, 90, 91, 92, 278, 312.
Jones, H. W., 43.
Jones, J. O., 254.
Kallabis, 388.
Karraker, P. E., 70, 71, 72.
Katutani, M., 310.
Keen, B. A., 260, 306.
Konekamp, A., 388.
Konig, 388.
Koshal, R. S., 258.
Kossovich, P., 462.
Kraebel, C. J., 9.
Krugel, C., 395.
Krüger, E., 357, 358, 362, 370.
Laatsch, W., 399.
Ladd, E. F., 100.
Lake States Forest Experiment Station, 205.
Latham, B., 247.
Lawes, J. B., 221, 222, 223, 224, 225, 226, 228.
Leather, J. W., 284, 287, 288, 289, 290, 291.
Lee, C. H., 6.
Leland, E. W., 123.
Liebig, J. von, 321.
Lowdermilk, W. C., 3, 4.
Luedecke, C., 244.
Lugner, I., 456, 457.
Lunt, H. A., 19, 23.
Lyon, T. L., 104, 106, 109, 112, 115, 120, 123, 127, 133, 178, 248, 250, 253, 270, 369, 372.
MacIntire, W. H., 152, 154, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 179, 180, 181, 182, 183, 184, 186, 187, 188, 189, 190, 191, 192, 193, 201.
Maluschitzki, N., 459.
Marriott, W., 247.
Martin, J. C. 204.
Maschaupt, J. G., 452.
Mawley, E., 231, 238.
Maxwell, W., 47, 48, 49.
McGeorge, W. T., 1.
McKaig, N., Jr., 150, 151, 151a.
Mellish, H., 247.
Mensching, J. E., 122, 125.
Miege, E., 487.
Miller, M. F., 86.
Miller, N. H. J., 239, 242, 243.
Minssen, 347.
Mollendorff, G., 320.
Mooers, C. A., 152, 153, 155, 158, 173, 179, 188.
Morgan, M. F., 14, 21, 24.
Mosier, J. G., 251.
Mukerji, J. N., 298.
Musgrave, G. W., 63, 64, 65, 67, 68.
Muther, A., 353.
Nath, B. V., 307.
Neal, O. R., 68, 69.
New York State Agricultural Experiment Station, 93, 94, 105. 107, 110, 113, 116, 121, 124, 126, 129, 130, 132, 134, 135.
Nicholson, H. H., 256.
Niklas, H., 382, 391.
North of Scotland College of Agriculture, 264, 265, 281.
Norton, R. A., 67.
Novikoff, V., 486.
Osugi, S., 313.
Palmer, V. J., 206.
Pantin, B., 256.
Pantoli, B., 440.
Parkes, J., 214.
Paterson, J. W., 262.
Peck, S. S., 55, 56, 57, 58.
Petermann, A., 401.
Pfaff, C., 398, 400.
Popov, V. P., 482.
Popp, M., 394.
Richards, E. H., 252.
Richards, L. A., 69.
Richter, G., 366, 367.
Richter, L., 463, 476.
Riesbol, H. S., 140.
Risler, E., 434, 435.
Robertson, I. M., 277.
Robinson, B., 168, 189, 190, 192, 193.
Robinson, G. W., 254, 269.
Roller, E. M., 151, 151a.

Ross, J. C., 489.
Rothamsted Experiment Station, 259, 261.
Ruprecht, R. W., 34, 35, 36, 37, 38, 40
Russell, E. J., 185, 241, 246, 252, 257, 273.
Russell, M. B., 69.
Rutkovsky, V. J., 483.
Sanborn, J. W., 85.
Sanders, K. B., 172, 177, 181, 187, 189.
Scheffler, H., 336, 337.
Scholz, H. F., 207.
Schropp, W., 382, 391.
Schwarz, 388.
Scott, R. H., 238.
Seelhorst, C. von, 350, 351, 352, 353, 354, 355, 361, 365.
Sen, J., 293.
Seyfert, F., 232.
Shaw, W. M., 160, 161, 162, 163, 164, 165, 167, 168, 172, 175, 180, 182, 183, 184, 187, 189, 190, 191, 192, 193, 201.
Shaw, W. N., 247.
Sherman, G. L., 140.
Shreve, F., 2.
Siegert, 368, 376, 379, 380, 383, 384, 388.
Slater, C. S., 137.
Smith, R. S., 59.
Staerk, 376, 379, 380, 383, 384.
Stauffer, R. S., 59.
Sterk, A. E., 443.
Stewart, A. B., 277.
Stockbridge, L., 79.
Stokes, W. E., 39.
Storer, F. H., 81, 235, 344.
Street, O. E., 11, 14.
Sturtevant, E. L., 73, 74, 75, 76, 77, 78.
Swanback, T. R., 11.
Tacke, 347.
Tamate, S., 312.
Taylor, G. H., 5.
Thayer, C. H., 83.
Thompson, Alice R., 52.
Thompson, W. R., 488.
Tornau, O., 393.
Treibich, 368.
Tret'iakov, I., 462.
Troubetzkoy, L., 461, 475.
Utz, E. J., 136.
Versluys, J., 447.
Walker, S. S., 31.
Warington, R., 102, 222, 223, 224, 225, 228, 237.
Warth, F. J., 297.
Welbel, B., 460, 463, 464, 466, 467, 468, 470, 472, 474, 476, 477, 479.
Welsh, H. D., 267, 268, 272, 276, 278, 280, 282.
Wery, G., 434, 435.
Wiley, H. W., 62, 411.
Willis, J. J., 233.
Willis, L. G., 154.
Wilms, 350.
Wilson, B. D., 117, 123.
Winkler, A., 465, 469, 471, 473, 478, 480.
Winnik, M., 308.
Wohltmann, F., 327, 335, 338.
Wollny, E., 326, 328, 329, 330, 333, 334, 339, 340, 341, 342, 343, 345, 401, 405, 408.
Wright, R. C., 209.
Yankovitch, L., 484.
Young, J. B., 159, 160, 161, 162, 163, 164, 166, 168, 173, 182, 184, 188, 193.
Zoller, R., 315, 318, 319.
Zunker, F., 389.

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE WHEN THIS PUBLICATION WAS LAST PRINTED

Secretary of Agriculture	HENRY A. WALLACE.
Under Secretary	CLAUDE W. WICKARD.
Assistant Secretary	GROVER B. HILL.
Director of Information	M. S. EISENHOWER.
Director of Extension Work	M. L. WILSON.
Director of Finance	W. A. JUMP.
Director of Personnel	ROY F. HENDRICKSON.
Director of Research	JAMES T. JARDINE.
Director of Marketing	MILO R. PERKINS.
Solicitor	MASTIN G. WHITE.
Land Use Coordinator	M. S. EISENHOWER.
Office of Plant and Operation	ARTHUR B. THATCHER, *Chief.*
Office of C. C. C. Activities	FRED W. MORRELL, *Chief.*
Office of Experiment Stations	JAMES T. JARDINE, *Chief.*
Office of Foreign Agricultural Relations	LESLIE A. WHEELER, *Director.*
Agricultural Adjustment Administration	R. M. EVANS, *Administrator.*
Bureau of Agricultural Chemistry and Engineering.	HENRY G. KNIGHT, *Chief.*
Bureau of Agricultural Economics	H. R. TOLLEY, *Chief.*
Agricultural Marketing Service	C. W. KITCHEN, *Chief.*
Bureau of Animal Industry	JOHN R. MOHLER, *Chief.*
Commodity Credit Corporation	CARL B. ROBBINS, *President.*
Commodity Exchange Administration	J. W. T. DUVEL, *Chief.*
Bureau of Dairy Industry	O. E. REED, *Chief.*
Bureau of Entomology and Plant Quarantine	LEE A. STRONG, *Chief.*
Farm Credit Administration	A. G. BLACK, *Governor.*
Farm Security Administration	W. W. ALEXANDER, *Administrator.*
Federal Crop Insurance Corporation	LEROY K. SMITH, *Manager.*
Federal Surplus Commodities Corporation	MILO R. PERKINS, *President.*
Food and Drug Administration	WALTER G. CAMPBELL, *Chief.*
Forest Service	EARLE H. CLAPP, *Acting Chief.*
Bureau of Home Economics	LOUISE STANLEY, *Chief.*
Library	CLARIBEL R. BARNETT, *Librarian.*
Division of Marketing and Marketing Agreements.	MILO R. PERKINS, *In Charge.*
Bureau of Plant Industry	E. C. AUCHTER, *Chief.*
Rural Electrification Administration	HARRY SLATTERY, *Administrator.*
Soil Conservation Service	H. H. BENNETT, *Chief.*
Weather Bureau	FRANCIS W. REICHELDERFER, *Chief.*

This publication is a contribution from

Soil Conservation Service	H. H. BENNETT, *Chief.*
Office of Research	M. L. NICHOLS, *Asst. Chief in Charge.*
Hydrologic Division	C. E. RAMSER, *Chief.*
Conservation Experiment Stations Division.	A. E. BRANDT, *Acting Chief.*

U. S. GOVERNMENT PRINTING OFFICE: 1940